

A233

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
22 January 2004 (22.01.2004)

PCT

(10) International Publication Number  
**WO 2004/007743 A2**

- (51) International Patent Classification<sup>7</sup>: **C12Q**
- (21) International Application Number: PCT/IB2003/003727
- (22) International Filing Date: 17 July 2003 (17.07.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
60/396,432 17 July 2002 (17.07.2002) US
- (71) Applicant: **COLEY PHARMACEUTICAL GMBH**  
[DE/DE]; Elisabeth-Selbert-Strasse 9, 40764 Langenfeld  
(DE).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **WAGNER, Her-  
mann** [DE/DE]; Kaagangerstrasse 36, 82279 Eching  
(DE). **KRETZSCHMAR, Hans** [DE/DE]; Nelkenweg  
5a, 82515 Wolfratshausen (DE). **SETHI, Shneh** [DE/DE];  
Platanen Strasse 56, 81377 Muenchen (DE).
- (74) Agent: **STEELE, Alan, W.**; Wolf, Greenfield & Sacks,  
P.C., 600 Atlantic Avenue, Boston, MA 02210 (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC,  
SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA,  
UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,  
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,  
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— without international search report and to be republished  
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

WO 2004/007743 A2

(54) Title: USE OF CPG NUCLEIC ACIDS IN PRION-DISEASE

(57) Abstract: Methods are provided which are useful in the treatment of prion diseases and other protein deposit diseases, includ-  
ing, for example, post-exposure prophylaxis against the development of iatrogenic Creutzfeldt-Jakob disease. The methods involve  
the use of immunostimulatory nucleic acids, including CpG nucleic acids.

- 1 -

## USE OF CpG NUCLEIC ACIDS IN PRION DISEASE

### Field of the Invention

The instant invention pertains to methods useful in the treatment of prion diseases, including, for example, post-exposure prophylaxis against the development of iatrogenic Creutzfeldt-Jakob disease. The methods involve the use of immunostimulatory nucleic acids.

### Background of the Invention

Prion diseases include a number of fatal, neurodegenerative diseases believed to be caused by aggregates of normal protein that is present in an abnormal conformation. The normal protein, prion protein, is usually present in the cell membrane of many tissues, particularly neuronal tissue. The abnormally conformed prion protein is believed to be directly involved in converting normally conformed prion protein into more of the abnormally conformed prion protein, which then self-assembles into aggregates that are damaging to neuronal tissue anatomy and function.

At least some of the prion diseases are transmissible. However, unlike bacteria, viruses, fungi, parasites, and other replicating pathogens, transmissible prions are simply proteins; they are transmissible without any accompanying nucleic acid. For reasons that are not yet fully understood, the abnormally conformed prion proteins normally do not induce an immune response. Thus, exposure of a healthy individual to abnormally conformed prion protein can initiate a prion disease that can go unchecked by the immune system.

Exposure to abnormally conformed prion protein thus represents a health risk to susceptible individuals. Such individuals include humans and non-humans, principally cows and sheep. Exposure can come about through contact with prion-diseased animal products through ingestion, iatrogenic or work-related exposure, transplantation, and administration of pharmaceutical preparations.

### Summary of the Invention

The instant invention is based in part on the unexpected discovery by the inventors that administration of immunostimulatory nucleic acid to a subject that is exposed to abnormally conformed prion protein is an effective treatment of prion disease. The immunostimulatory nucleic acid effectively delayed and even prevented disease in animals

- 2 -

administered very large doses of prion-diseased brain homogenates that uniformly caused fatal disease in all untreated control animals similarly exposed to prion-diseased brain homogenates.

The immunostimulatory nucleic acids are useful in the treatment of prion diseases, including Creutzfeldt-Jakob disease (CJD), bovine spongiform encephalopathy (BSE), and scrapie. The methods of the invention are also useful for the study of these diseases, for instance, in animal models. The methods will also be useful for developing an understanding of how prion proteins normally fail to elicit an immune response and how an immune response can be elicited and used to treat prion protein disease.

The immunostimulatory nucleic acids are also useful in the treatment of other neurologic diseases involving abnormal protein deposits or aggregates. Such diseases include Alzheimer's disease, which involves deposits of amyloid. The main component of amyloid plaques is amyloid- $\beta$  peptide (A $\beta$ ), a fibrillar 40-42 amino acid peptide that accumulates extracellularly and causes neuronal death.

In one aspect the invention provides a method for treating a prion disease in a subject. The method involves administering to a subject having or at risk of developing a prion disease a CpG nucleic acid in an effective amount to treat the prion disease. In one embodiment the administering follows exposure of the subject to a prion protein that is associated with a prion disease. In one embodiment the prion disease is a transmissible spongiform encephalopathy (TSE). In one embodiment the subject is a human.

In various preferred embodiments, the prion disease is scrapie, BSE, or a form of CJD. The CJD in one embodiment is iatrogenic CJD (iCJD). In another embodiment the CJD is variant CJD (vCJD). In yet another embodiment the CJD is sporadic CJD.

In one aspect the invention provides a method for inducing an immune response to a prion protein. The method according to this aspect of the invention involves the steps of contacting an antigen-presenting cell (APC) with a prion protein and contacting the APC with a CpG nucleic acid in an effective amount to induce an immune response to the prion protein. In one embodiment the immune response occurs in vivo. In one embodiment the immune response occurs in vitro. In all embodiments according to this aspect of the invention the APC is preferably chosen from a B cell, a dendritic cell, a macrophage, and a monocyte. In one preferred embodiment the APC is a dendritic cell.

- 3 -

Also according to this aspect of the invention, in one embodiment the APC expresses a Toll-like receptor (TLR) that signals in response to the CpG nucleic acid. It has recently been reported that CpG nucleic acid can specifically induce a particular TLR, designated TLR9, to signal. Accordingly, in one embodiment the TLR is TLR9.

In one embodiment the prion protein includes prion protein:scrapie form (PrP<sup>Sc</sup>). In another embodiment the prion protein includes a fragment of PrP<sup>Sc</sup> lacking at least the amino terminus of full-length PrP<sup>Sc</sup>. In yet another embodiment the prion protein includes a derivative of PrP<sup>Sc</sup> or a derivative of a fragment of PrP<sup>Sc</sup> lacking at least the amino terminus of full-length PrP<sup>Sc</sup>.

In one embodiment the prion protein is prion protein:scrapie form (PrP<sup>Sc</sup>). In another embodiment the prion protein is a fragment of PrP<sup>Sc</sup> lacking at least the amino terminus of full-length PrP<sup>Sc</sup>. In yet another embodiment the prion protein is a derivative of PrP<sup>Sc</sup> or a derivative of a fragment of PrP<sup>Sc</sup> lacking at least the amino terminus of full-length PrP<sup>Sc</sup>.

It has been reported that certain CpG nucleic acids are more effective in one species than in another. Accordingly, in preferred embodiments the CpG nucleic acid is optimized for use in a species of the subject.

It has also been reported that CpG nucleic acids appear to fall into different classes based on certain structural features as well as their function. At least three classes are believed to exist, denoted Class A, Class B, and Class C. In one embodiment the CpG nucleic acid is a class B CpG nucleic acid. In one embodiment the CpG nucleic acid is a class A CpG nucleic acid. In one embodiment the CpG nucleic acid is a class C CpG nucleic acid.

These and other features of the invention are described in greater detail in connection with the detailed description of the invention.

### **Detailed Description of the Invention**

A major step in the study of prions and the diseases that they cause was the discovery and purification of a protein designated prion protein (PrP). Bolton et al. (1982) *Science* 218:1309-11; Prusiner SB et al. (1982) *Biochemistry* 21:6942-50; McKinley MP et al. (1983) *Cell* 35:57-62. Complete prion protein-encoding genes have since been cloned, sequenced and expressed in transgenic animals. PrP<sup>C</sup> is encoded by a single-copy host gene (Basler K et

- 4 -

al. (1986) *Cell* 46:417-28) and is normally found at the outer surface of neurons. Prion diseases are accompanied by the conversion of PrP<sup>C</sup> into a modified form called PrP<sup>Sc</sup>.

The scrapie isoform of the prion protein (PrP<sup>Sc</sup>) is thought necessary for both the transmission and pathogenesis of the transmissible neurodegenerative diseases of animals and humans. See Prusiner SB (1991) *Science* 252:1515-22. The most common prion diseases of animals are scrapie of sheep and goats and bovine spongiform encephalopathy (BSE; "mad cow disease") of cattle. Wilesmith J et al. (1991) *Microbiol Immunol* 172:21-38. Four prion diseases of humans have been identified: (1) kuru, (2) Creutzfeldt-Jakob disease (CJD), (3) Gerstmann-Sträussler-Scheinker (GSS) syndrome, and (4) fatal familial insomnia. Gajdusek DC (1977) *Science* 197:943-60; Medori et al. (1992) *N Engl J Med* 326:444-9.

Most CJD cases are sporadic, but about 10-15% are inherited as autosomal dominant disorders that are caused by mutations in the human PrP gene. Hsiao et al. (1990) *Neurology* 40:1820-7; Goldfarb et al. (1992) *Science* 258:806-8. Iatrogenic CJD has been caused by human growth hormone derived from cadaveric pituitaries as well as dura mater grafts. Brown et al. (1992) *Lancet* 340:24-7. Despite numerous attempts to link CJD to an infectious source such as the consumption of meat from scrapie-infected sheep, none has been identified to date (Harries-Jones et al. (1988) *J Neurol Neurosurg Psychiatry* 51:1113-9) except in cases of iatrogenically induced disease. On the other hand, kuru, which for many decades devastated the Fore and neighboring tribes of the New Guinea highlands, is believed to have been spread by infection during ritualistic cannibalism.

The major component of purified infectious prions, designated PrP 27-30, is the proteinase K resistant core of a larger native protein PrP<sup>Sc</sup> which is the disease causing form of the ubiquitous cellular protein PrP<sup>C</sup>. PrP<sup>Sc</sup> is found only in scrapie infected cells, whereas PrP<sup>C</sup> is present in both infected and uninfected cells implicating PrP<sup>Sc</sup> as the major, if not the sole, component of infectious prion particles. Since both PrP<sup>C</sup> and PrP<sup>Sc</sup> are encoded by the same single copy gene, great effort has been directed toward unraveling the mechanism by which PrP<sup>Sc</sup> is derived from PrP<sup>C</sup>. Central to this goal has been the characterization of physical and chemical differences between these two molecules. Properties distinguishing PrP<sup>Sc</sup> from PrP<sup>C</sup> include low solubility (Meyer RK et al. (1986) *Proc Natl Acad Sci USA* 83:2310-4), poor antigenicity (Kascsak RJ et al. (1987) *J Virol* 61:3688-93; Serban D et al. (1990) *Neurology* 40:110-7), protease resistance (Oesch B et al. (1985) *Cell* 40:735-46), and polymerization of PrP 27-30 into rod-shaped aggregates which are very similar, on the

- 5 -

ultrastructural and histochemical levels, to the PrP amyloid plaques seen in scrapie-diseased brains (Prusiner SB et al. (1983) *Cell* 35(2 Pt 1):349-58). By using proteinase K it is possible to denature PrP<sup>C</sup> but not PrP<sup>Sc</sup>. To date, attempts to identify any post-translational chemical modifications in PrP<sup>C</sup> that lead to its conversion to PrP<sup>Sc</sup> have proven fruitless. Stahl N et al. (1993) *Biochemistry* 31:5043-53. Consequently, it has been proposed that PrP<sup>C</sup> and PrP<sup>Sc</sup> are in fact conformational isomers of the same molecule.

Conformational description of PrP using conventional techniques has been hindered by problems of solubility and the difficulty in producing sufficient quantities of pure protein. However, PrP<sup>C</sup> and PrP<sup>Sc</sup> are conformationally distinct. Theoretical calculations based upon the amino acid sequences of PrPs from several species have predicted four putative helical motifs in the molecule. Experimental spectroscopic data would indicate that in PrP<sup>C</sup> these regions adopt alpha-helical arrangements, with virtually no beta-sheet. Pan et al. (1993) *Proc Natl Acad Sci USA* 90:10962-6). In dramatic contrast, in the same study it was found that PrP<sup>Sc</sup> and PrP 27-30 possess significant beta-sheet content, which is typical of amyloid proteins. Moreover, studies with extended synthetic peptides, corresponding to PrP amino acid residues 90-145, have demonstrated that these truncated molecules may be converted to either alpha-helical or beta-sheet structures by altering their solution conditions. The transition of PrP<sup>C</sup> to PrP<sup>Sc</sup> requires the adoption of beta-sheet structure by regions that were previously alpha-helical.

In general, scrapie infection fails to produce an immune response, with host organisms being tolerant to PrP<sup>Sc</sup> from the same species. Polyclonal anti-PrP antibodies have been raised in rabbits following immunization with large amounts of Syrian hamster PrP 27-30. Bendheim PE et al. (1985) *Proc Natl Acad Sci USA* 82:997-1001; Bode L et al. (1985) *J Gen Virol* 66:2471-8. Similarly, a handful of anti-PrP monoclonal antibodies have been produced in mice. Kascsak RJ et al. (1987) *J Virol* 61:3688-93; Barry RA et al. (1986) *J Infect Dis* 154:518-21. These antibodies are able to recognize native PrP<sup>C</sup> and denatured PrP<sup>Sc</sup> from both Syrian hamsters and humans equally well, but do not bind to murine PrP. Unsurprisingly, the epitopes of these antibodies were mapped to regions of sequence containing amino acid differences between Syrian hamster and murine PrP. Rogers et al. (1991) *J Immunol* 147:3568-74.

The DNA sequence of the human, sheep and cow PrP genes have been determined, allowing, in each case, the prediction of the complete amino acid sequence of their respective

- 6 -

PrP proteins. The normal amino acid sequence which occurs in the vast majority of individuals is referred to as the wild-type PrP sequence. This wild-type sequence is subject to certain characteristic polymorphic variations. In the case of human PrP, two polymorphic amino acids occur at residues 129 (Met/Val) and 219 (Glu/Lys). Sheep PrP has two amino acid polymorphisms at residues 136 and 171, while bovine PrP has either five or six repeats of an eight amino acid motif sequence (octarepeats) in the amino terminal region of the mature prion protein. While none of these polymorphisms are of themselves pathogenic, they appear to influence prion diseases. Distinct from these normal variations of the wild-type PrP proteins, certain mutations of the human PrP gene which alter either specific amino acid residues of PrP or the number of octarepeats have been identified which segregate with inherited human prion diseases.

For example, sequences of chicken, bovine, sheep, rat, and mouse PrP genes are disclosed and published within Gabriel JM et al. (1992) *Proc Natl Acad Sci USA* 89:9097-101. A sequence for a PrP gene of Syrian hamster is published in Basler K et al. (1986) *Cell* 46:417-28. A PrP gene of sheep is published by Goldmann W et al. (1990) *Proc Natl Acad Sci USA* 87:2476-80. A gene sequence for bovine PrP is published in Goldmann W et al. (1991) *J Gen Virol* 72:201-4. A sequence for chicken PrP gene is published in Harris DA et al. (1991) *Proc Natl Acad Sci USA* 88:7664-8. A PrP gene sequence for mink is published in Kretzschmar HA et al. (1992) *J Gen Virol* 73:2757-61. A human PrP gene sequence is published in Kretzschmar HA et al. (1986) *DNA* 5:315-24. A PrP gene sequence for mouse is published in Loch C et al. (1986) *Proc Natl Acad Sci USA* 83:6372-6. A PrP gene sequence for sheep is published in Westaway D et al. (1994) *Genes Dev* 8:959-69. These publications are all incorporated herein by reference to disclose and describe the PrP gene and PrP amino acid sequences.

#### Human PrP cDNA (SEQ ID NO:1; GenBank Accession No. M13899)

|          |           |           |            |            |            |     |
|----------|-----------|-----------|------------|------------|------------|-----|
| cggcgccg | agcttcct  | ctctcacga | ccgaggcaga | gcagtcatta | tggcgaacct | 60  |
| tggctgct | atgctggt  | tctttgtgg | cacatggagt | gacctggg   | tctgcaagaa | 120 |
| gcgcccga | cctggagg  | ggaacactg | gggcagccga | taccgggg   | agggcagccc | 180 |
| tggaggca | cgtaccac  | ctcagggcg | tggtggctg  | gggcagcct  | atggtggtg  | 240 |
| ctggggcg | cctcatgg  | gtggctggg | gcagcccat  | ggtggtgg   | ggggacagcc | 300 |
| tcattggt | ggctgggt  | aaggaggtg | caccacagt  | cagtggaac  | agccgagtaa | 360 |
| gcaaaaac | aacatga   | acatggctg | tgctgcagc  | gctggggc   | tggtggggg  | 420 |
| cctggcgc | tacatgct  | gaagtgcc  | gagcaggcc  | atcatacat  | tcggcagtga | 480 |
| ctatgagg | cgttactat | gtgaaaac  | gcaccgttac | cccaacca   | tgtactacag | 540 |
| gcccattg | gagtacag  | accagaaca | ctttgtgc   | gactgcgtc  | atatcacaat | 600 |
| caagcagc | acggtcac  | caaccacaa | gggggaga   | ttcaccgag  | ccgacgttaa | 660 |
| gatgatgg | cgcgtggt  | agcagatgt | tatcaccag  | tacgagagg  | aatctcaggc | 720 |

- 7 -

|             |             |            |             |             |            |      |
|-------------|-------------|------------|-------------|-------------|------------|------|
| ctattaccag  | agaggatcga  | gcatggctct | cttctcctct  | ccacctgtga  | tcctcctgat | 780  |
| ctctttcctc  | atcttctga   | tagtgggatg | aggaaggtct  | tcctgttttc  | accatctttc | 840  |
| taatcttttt  | ccagcttgag  | ggaggcggta | tccacctgca  | gcccttttag  | tgggtggtgc | 900  |
| tcactctttc  | ttctctcttt  | gtcccggata | ggctaataca  | tacccttggc  | actgatgggc | 960  |
| actggaaaac  | atagagtaga  | cctgagatgc | tggtcaagcc  | ccctttgatt  | gagttcatca | 1020 |
| tgagccgttg  | ctaagccag   | gccagtaaaa | gtataacagc  | aaataacccat | tggttaatct | 1080 |
| ggacttattt  | ttggacttag  | tgcaacaggt | tgaggctaaa  | acaaatctca  | gaacagtctg | 1140 |
| aaataccctt  | gcctgggatac | ctctggctcc | ttcagcagct  | agagctcagt  | atactaatac | 1200 |
| cctatcttag  | tagagatttc  | atagctattt | agagatattt  | tccattttta  | gaaaacccga | 1260 |
| caacatttct  | gccaggtttg  | ttaggaggcc | acatgatact  | tattcaaaaa  | aatcctagag | 1320 |
| attcttaggt  | cttgggatgc  | aggctcagcc | cgtggagca   | tgagctctgt  | gtgtaccgag | 1380 |
| aactgggggtg | atgttttact  | tttcacagta | tgggctacac  | agcagctgtt  | caacaagagt | 1440 |
| aaatatgtgc  | acaacactga  | acctctggct | agaggacata  | ttcacagtga  | acataactgt | 1500 |
| aacatatatg  | aaaggcttct  | gggacttgaa | atcaaagtgt  | tgggaatggg  | gcccttggag | 1560 |
| gcaacctccc  | attttagatg  | tttaaaggac | cctatatgtg  | gcattccttt  | ctttaaacta | 1620 |
| taggtaatta  | aggcagctga  | aaagtaaat  | gccttctaga  | cactgaaggc  | aaatctcctt | 1680 |
| tgtccattta  | cctggaaacc  | agaatgattt | tgacatacag  | gagagctgca  | gttgtgaaag | 1740 |
| caccatcatc  | atagaggatg  | atgtaattaa | aaaatgggtca | gtgtgcaaag  | aaaagaactg | 1800 |
| cttgcatttc  | tttattcttg  | ttcacaattt | gtcaaaaacc  | agaattaggt  | caagttcata | 1860 |
| gtttctgtaa  | ttggcttttg  | aatcaaagaa | tagggagaca  | atctaaaaaa  | tatcttaggt | 1920 |
| tgagatgac   | agaaatatga  | ttgatttgaa | gtggaaaaag  | aaattctgtt  | aatgttaatt | 1980 |
| aaagtaaaat  | tattccctga  | attgtttgat | attgtcacct  | agcagatatg  | tattactttt | 2040 |
| ctgcaatggt  | attattggct  | tgcactttgt | gagtatctat  | gtaaaaatat  | atatgtatat | 2100 |
| aaaatatata  | ttgcatagga  | cagacttagg | agttttgttt  | agagcagtta  | acatctgaag | 2160 |
| tgtctaagtc  | attaactttt  | gtaaggctac | gaatacttaa  | tatgtgggaa  | acccttttgc | 2220 |
| gtggctctta  | ggcttacaat  | gtgcactgaa | tcgtttcatg  | taagaatcca  | aagtggacac | 2280 |
| cattaacagg  | cttttgaaat  | atgcatgtac | tttatatttt  | ctatatattgt | aactttgcat | 2340 |
| gttcttgttt  | tgttatataa  | aaaaattgta | aatgtttaat  | atctgactga  | aattaaacga | 2400 |
| gcgaagatga  | gcacc       |            |             |             |            | 2415 |

## Mink PrP genomic DNA (SEQ ID NO:2; GenBank Accession No. S46825)

|             |            |            |             |            |            |      |
|-------------|------------|------------|-------------|------------|------------|------|
| tcattttggt  | ttgttttgtt | ttgtttgcag | ataagccatc  | atggtgaaaa | gccacatagg | 60   |
| cagctggctc  | ctggttctct | ttgtggccac | atggagtgc   | attggcttct | gcaagaagcg | 120  |
| gccaaagcct  | ggaggaggct | ggaacactgg | ggggagccga  | taccagggc  | agggcagtc  | 180  |
| tgagggaac   | cgctaccac  | cccagggtgg | tggcggctgg  | ggccagcccc | acgggggtgg | 240  |
| ctggggacag  | cccacggggg | gtggctgggg | tcagccccc   | gggggtggct | ggggacagcc | 300  |
| gcatgggtggc | ggtggctggg | gtcaaggtgg | tgggagccac  | ggtcagtggt | gcaagcccag | 360  |
| taagcccaaa  | accaacatga | agcatgtggc | gggagccgca  | gcagccgggg | cggtcgtggg | 420  |
| gggcctgggc  | ggctacatgc | tggggagcgc | catgagcagg  | cccctcattc | attttggcaa | 480  |
| cgactatgag  | gaccgtact  | accgtgagaa | catgtaccgc  | tacccaacc  | aagtgtacta | 540  |
| caagccgggtg | gatcagtaca | gcaaccagaa | caacttcgtg  | catgactgcg | tcaacatcac | 600  |
| ggtcaagcag  | cacacggtga | ccaccaccac | caagggcgag  | aacttcacgg | agaccgacat | 660  |
| gaagatcatg  | gagcgcgtgg | tggagcagat | gtgtgtcacc  | cagtaccagc | gagagtcgca | 720  |
| ggcttactac  | cagagggggg | cgagcgccat | cctcttctcg  | ccccctccc  | tgatcctcct | 780  |
| catctcactg  | ctcattctcc | tgatagtggg | atgaggatgg  | ccttcccatt | ctctccatcg | 840  |
| tcttcacctt  | ttacagggtg | ggggaggggg | tgtctacct   | cagccctgta | gtgggtgggt | 900  |
| ctcattcctg  | cttctcttta | tcacccatag | gctaataccc  | ttggccctga | tggccctggg | 960  |
| aaatgtagag  | cagacccagg | atgctattta | ttcaagcccc  | catgtgttgg | agtccttcag | 1020 |
| gggccaatgc  | tagtgcaggg | ctgagaataa | cagcaaatca  | tcattggttg | acctagggct | 1080 |
| gcttttttgt  | tgtgtgtgtc | tagtgcagct | gaccgaggct  | aaaacaattc | tcaaaacagt | 1140 |
| tttcaaatac  | ctttgccttg | aaacctctgg | ctcctgtctg  | agctagagct | cagtacatta | 1200 |
| atgtcccatc  | ttagccgtgt | cttcatagca | acttggggaa  | gtttttctcc | ccactctaaa | 1260 |
| agaacgcgat  | tgcacttccc | tgtgcaaaga | acatttctgc  | caaatttgaa | aggaggccac | 1320 |
| atgatattca  | ttcaaaaagc | aaaactagaa | accctttgct  | cttggacgca | agcccgccct | 1380 |
| gctaggagca  | ccaaactggg | gcgatgggtt | ctactctgcg  | gcgtgggcta | tggcgacgcc | 1440 |
| gaggtgtcca  | gcgtaaatat | tgatgcgacg | ctagacctag  | gcagaggatg | tttgcacagg | 1500 |
| gaatgaacat  | aatcaacagt | gcgaaaatgc | tacaaaaaat  | cccacactgg | ggagcagtgt | 1560 |
| ccttggaggc  | aagttttttt | ccttttggga | catttaaaagc | ccctatatgt | ggcattcctt | 1620 |



- 8 -

|             |             |            |            |             |            |      |
|-------------|-------------|------------|------------|-------------|------------|------|
| tctttcgtaa  | cctaaactat  | agataattaa | ggcagttaaa | aattgaactt  | ccttccaggc | 1680 |
| cccaagagca  | aatctttgtt  | cacttacctg | gaaaccagaa | tgattttgac  | acagaggaag | 1740 |
| gtgcagctgt  | taaaataacc  | ctcatcctag | aagattgcat | catggagaaa  | acgatccgta | 1800 |
| gacaaaaatg  | atcgcatctt  | ttcattgctg | tctcgtaatt | gacagaaacc  | agaattatgt | 1860 |
| caagtcctag  | tttctataat  | cagcttttga | atcaaagaat | ggaagtccat  | ccaaaaaaaa | 1920 |
| aaaagaaata  | ccttaggtca  | cccatgacag | aaatacccat | tcagggttaga | aaaaaggaat | 1980 |
| tctgttaact  | gttatattaag | taaggcaaaa | ttattgtccg | gattgttcga  | tatcatcagc | 2040 |
| tagcagataa  | attagcattc  | tgcaatgttc | cgggcttgca | ctgtgcgggt  | atttgatgtt | 2100 |
| aaaaaaaaatt | attatatata  | ttgtgtatga | caaacttaga | agtttttgct  | agaggagtta | 2160 |
| acatctgata  | tatctaatagc | accaccagtt | ttggaaggta | ctaaataactt | aatatgtaga | 2220 |
| aatccttttg  | cgtggctctc  | aggcttacac | gtgcactgaa | tagttttgta  | tgatagagcc | 2280 |
| ctgtgggtct  | tcgaaatatg  | catgtacttt | atattttcta | tatttgtaac  | tgggcatgta | 2340 |
| cttgataaaa  | aaatgtataa  | acattcgaac | tcttgactag | aattaaacag  | gaactgagtg | 2400 |
| tgtcccatgt  | gtttgcagtg  | acattcacca | cgcacccctg | tgttgg      |            | 2446 |

## Syrian Hamster PrP cDNA (SEQ ID NO:3; GenBank Accession No. M14054)

|             |             |            |            |             |            |     |
|-------------|-------------|------------|------------|-------------|------------|-----|
| tcgaaaaatct | ccctcttttag | caatttcttg | ctcctagagt | ttcagcaatt  | gctttctcgc | 60  |
| tccatttaggc | aacctttcat  | tttctcacct | tcccattat  | gtaacgggag  | caatgggttc | 120 |
| tggaccagtc  | ttccattaaa  | gatgattttt | atagtcggtg | agcgccgtca  | gggagtgatg | 180 |
| acacctgggg  | gcggtttaaa  | ccgtacaatc | ccttaaacca | gtctggagcg  | gtgactcatt | 240 |
| ttcccaggga  | gaagtggcgc  | ggccattggt | gagcacgacg | caagccccgc  | cccaccagc  | 300 |
| ccggccccgc  | cctgctacc   | ctcctgactc | actgccccgc | ccgctcccc   | gcggcgtccg | 360 |
| agcagcagac  | cgagaaggca  | catcgagtcc | actcgtcgcg | tcggtggcag  | gtaagcggct | 420 |
| tctgaagcct  | ggccccggga  | aggggtgctg | agccaggcct | cggtaagcct  | tcggcttccc | 480 |
| agagccaagc  | ccggcttact  | ccggctctcg | gggcgctgag | gccgcggggc  | tgagggttag | 540 |
| tctggctggg  | aggtgaccgc  | gcaccgcgag | ccgcgcgtct | ccttgaggga  | ccgaacccca | 600 |
| ggagaggcca  | ggagccatcc  | cttcctcccg | agccccgctc | acccccagag  | tcgctcgggg | 660 |
| atgggggatg  | ggggatgggg  | tggcatcttt | tgactgtcgt | ttgctgtttt  | cttctctctt | 720 |
| tgtaatagct  | acagcgaaca  | taattttacc | cagggttcca | ccgtgggtctc | gtccgctctc | 780 |
| ggcatctctc  | agtccagtac  | atacccaagg |            |             |            | 810 |

## Sheep PrP cDNA (SEQ ID NO:4; GenBank Accession No. AJ223072)

|             |            |             |             |            |            |      |
|-------------|------------|-------------|-------------|------------|------------|------|
| atggtgaaaa  | gccacatagg | cagttggatc  | ctggttctct  | ttgtggccat | gtggagtga  | 60   |
| gtgggcctct  | gcaagaagcg | acaaaaacct  | ggcggaggat  | ggaacactgg | ggggagccga | 120  |
| taccggggac  | agggcagtc  | tggaggcaac  | cgctatccac  | ctcagggagg | gggtggctgg | 180  |
| ggtcagcccc  | atggaggtgg | ctggggccaa  | cctcatggag  | gtggctgggg | tcagccccat | 240  |
| gggtggtggct | ggggacagcc | acatggtggt  | ggaggctggg  | gtcaagggtg | tagccacagt | 300  |
| cagtggaaaca | agcccagtaa | gccccaaacc  | aacatgaagc  | atgtggcagg | agctgctgca | 360  |
| gctggagcag  | tggtaggggg | ccttgggtgg  | tacatgtcgg  | gaagtgccat | gagcaggcct | 420  |
| cttatacatt  | ttggcaatga | ctatgaggac  | cgttactatc  | gtgaaaacat | gtaccgttac | 480  |
| cccaaccaag  | tgtactacag | accagtggat  | cagtatagta  | accagaacaa | ctttgtgcat | 540  |
| gactgtgtca  | acatcacagt | caagcaacac  | acagtcacca  | ccaccaccaa | gggggagaac | 600  |
| ttcaccgaaa  | ctgacatcaa | gataatggag  | cgagtgggtg  | agcaaagtgt | catcaccag  | 660  |
| taccagagag  | aatcccaggc | ttattaccaa  | aggggggcaa  | gtgtgatcct | cttttcttcc | 720  |
| cctcctgtga  | tcctcctcat | ctctttcctc  | atttttctca  | tagtaggata | ggggcaacct | 780  |
| tcctgttttc  | attatcttct | taatctttgc  | cagggtgggg  | gagggagtgt | ctacctgcag | 840  |
| cctgtagtgt  | gtggtgtctc | atttcttgtc  | ctctcttgt   | tacctgtata | ataataacct | 900  |
| tggcgcttac  | agcactggga | aatgacaagc  | agacatgaga  | tgctgtttat | tcaagtccca | 960  |
| ttagctcagt  | attctaattg | cccatcttag  | cagtgtttt   | gtagcaattt | tctcatttgt | 1020 |
| ttcaagaaca  | cctgactaca | tttcccttgg  | ggaatagcat  | ttctgccaag | tctggaagga | 1080 |
| ggccacataa  | tattcattca | aaaaaacaaa  | actggaaatc  | cttagttcat | agaccagggt | 1140 |
| tccaccctgt  | tgagagcatg | tgtcctgtgt  | ctgcagagaa  | ctataaagga | tattctgcat | 1200 |
| tttgacaggt  | acatttgtag | gtaacacagc  | catctattgc  | atcaagaatg | gatattcatg | 1260 |
| caacctttga  | cttatgggca | gaggacatct  | tcacaaggaa  | tgaacataat | acaaaggctt | 1320 |
| ctgagactaa  | aaaattccaa | catatgggaag | aggtgccctt  | gggtggcagc | ttccattttg | 1380 |
| tatgtttaag  | acacttcaag | tgatatctct  | ttcttttagta | acataaagta | tagataatta | 1440 |
| aggtagctta  | attaaactac | cttctagaca  | ctgagagcaa  | atctgttgtt | tatctggaac | 1500 |

- 9 -

```

ccaggatgat tttgacattg cttagggatg tgagagttgg actgtaaaga aagctgagtg 1560
ctgaagaggt catgcttttg aactatagtg ttggagaaaa ctcttgagag tcccttgagc 1620
tgaaaggaga tcagtcctga atattcattg gaaggactga tgctgaagct gaaactccaa 1680
tactttgggt acctgatggg aagaactgaa ggcaggaggg atgctaggaa agactgaagg 1740
caggaggaga aggggacgac agaggatgag atggctagat ggcacatgag actcaatgga 1800
catgagctta agtaaaactcc aggagttggc aatggacagg gagacctggc gtcctgcagt 1860
ccatgggtgtc gcagagtcgg acacgattga gtgactaaat tgaggtgacc cagatttaac 1920
atagagaatg cagatacaaaa actcatattc atttgattga atcttttcct gaaccagtgc 1980
tagtgttgga ctggttaaggg tataacagca tatatagggt atgtgatgaa gagatagtg 2040
acatgaaata tgtgcatttc tttattgctg tcttataatt gtcaaaaaag aaaattaggt 2100
ccttggtttc tgtaaaattg acttgaatca aaaggagggc atttaaagaa ataaattaga 2160
gatgataaata atctgatcca ttcagagtag aaaaagaaat tccattactg ttattaaaga 2220
aggtaaaatt attccctgaa ttgttcaata ttgtcaccta gcagatagac actattctgt 2280
actgttttta ctagcttgca ccttggtgta tccatgtaa aaacatattt gcatatgaca 2340
aactttttct gttagagcaa ttaacatctg aaccacctaa tgcattacct gtttttgtaa 2400
ggtacttttt gtaaggtaact aaggagatgt ggggttaatc cctagggtcag gtaaatcccc 2460
tagaggaaga aatggcaacc cactccagta ttcttgccag gaaaatccag tgggcagagg 2520
agcctggcag ggtacagtct aagagcatgg ggttgcaaaag agtgagacaa gacttgagct 2580
actgaacaat aaggacaata aatgctgggt cggctaaaag gttcattagg tttttttct 2640
gtaaagatggc tctagttagta cttgtcttta tcttcattcg aaacaatttt gttagattgt 2700
atgtgacagc tcttgatca gcatgcattt gaaaaaaaca tcacaattgg taaatttttg 2760
tatagccatc ttactattga agatggaaga aaagaagcaa aattttcagc atatcatgct 2820
gtacttattt caagaaagat aaccaaagt caaaaatgta tttgtgaagt gtatggagaa 2880
ggggtgcaa ctgatcaagc ttgtcaaagt agtttgtaa gtttcgtgct ggagatttct 2940
tattggacga tgctccacag ttggatatac cagttgaagt tgatagtgat caaattgaga 3000
tattgagaat aatcgatgtt ataccacgag ggagatagct gacatactca aaatatccaa 3060
atagaacctt gaaaaccatt tgcaccatct cagttatgtt aatcactttg atgtttgagt 3120
tccacataag caaaaaaaca acaaaaaaaa aaaatacaac cttgaccata tttgcgcagt 3180
cagttctcta ctgaaatgat tgaaaacact ttgtttttaa aaacagattt tgattaacag 3240
tggttacgat acaataacgt agatggaaga aattgtaggg tgagcaaaat gaaccacacc 3300
accaaaaggc agtcttcctc taaagaagat gtgtgtatgg tgggattgga aagtaatcct 3360
ctattatgga ttcttctgga aaacactgct cctaattaga ccaactgaaa acagcactca 3420
acgaaaagca tccagaatta gtcaatagaa aacataatct tccatcagga taacgcaaga 3480
ctacatattt ctttgatgac ccagcatggc cggattttct gattcatctg ttgtttgagt 3540
acgttgcatc tttggatttt ttcatttat tttagtctac aaaattatca taatggaaaa 3600
aatttccatt ccctggaaga tgtaaagtgc atctggaaaa tttctttgct caaaaagata 3660
aaaagttttg tgaacacaga attatgacgt tgcctgaaaa atggcagaag gtatgggaac 3720
aaaagagtga ctatgttggt ttgtaaaagt cttagtgaat atgaaaaatg tgtcttttat 3780
ttttatttaa acaccaaagg cacattttag caaccataa ctgaatctaa aggaaactct 3840
tctgtgtgtt gtccttacag tgtgcactga tagtttgtat aagaatccag agtgatattt 3900
gaaatcgca tgtgcttata tttttatat ttgtaacttt gcattgtact gttttgtgtt 3960
aaaagtttat aaatatttaa tatctgacta aaattaaaac ggagctaaaa ggagtatctt 4020

```

## Bovine PrP cDNA (SEQ ID NO:5; GenBank Accession No. X55882)

```

atgggtgaaaa gccacatagg cagttggatc ctggttctct ttgtggccat gtggagtgac 60
gtgggcctct gcaagaagcg accaaaacct ggaggaggat ggaacactgg ggggagccga 120
taccaggagc agggcagtc tggaggcaac cgttatccac ctgaggagg ggggtgctgg 180
ggtcagcccc atggaggtgg ctggggccag gctcatggag gtggctgggg ccagcctcat 240
ggaggtggct ggggtcagcc ccatggtggg ccttggggac agccacatgg tgggtgaggc 300
tggggtcaag gtggtaccca cggatcaatg aacaaacca gtaagccaaa aaccaacatg 360
aagcatgtgg caggagctgc tgcagctgga gcagtggtag ggggccttgg tggctacatg 420
ctgggaagtg ccatgagcag gcctcttata cathttggca gtgactatga ggaccgttac 480
tatcgtgaaa acatgcaccg ttacccaac caagtgtact acaggccagt ggatcagtat 540
agtaaccaga acaactttgt gcatgactgt gtcaacatca cagtcaagga acacacagtc 600
accaccacca ccaaggggga gaacttcacc gaaactgaca tcaagatgat ggagcgagtg 660
ctggagcaaaa tgtgcattac ccagtaccag agagaatccc aggcctatta ccaacgaggg 720
gcaagtgtga tctctctctc tccccctcct gtgatcctcc tcatctcttt cctcattttt 780
ctcatagtag gatag 795

```

- 10 -

## Chicken PrP cDNA (SEQ ID NO:6; GenBank Accession No. M61145)

|            |            |            |            |            |            |      |
|------------|------------|------------|------------|------------|------------|------|
| gaattccctc | ggcagccagc | tctccctct  | cgctatttat | tcctttctcc | ccccctacg  | 60   |
| ctggatctgg | atcatctcaa | gccgagcgg  | gacggcttct | tggatcgctc | atacataaat | 120  |
| atctgtgagt | cagaggaagc | aaccaccgac | cccaagacct | caccccgagc | catggctagg | 180  |
| ctctcacca  | cctgtgcct  | gctggccctg | ctgctcgccg | cctgcaccga | cgctgcctc  | 240  |
| tccaagaagg | gcaaaggcaa | accagtggt  | gggggttggg | gcgccgggag | ccatcgccag | 300  |
| cccagctacc | cccgccagcc | gggctaccct | cataaccag  | ggtaccccca | taaccaggg  | 360  |
| tacccccaca | accctggcta | tcccataac  | cccggctacc | cccagaacct | tggctacccc | 420  |
| cataaccag  | gttaccag   | ctggggta   | ggctacaacc | catccagcgg | aggaagtac  | 480  |
| cacaaccaga | agccatggaa | acccccaaa  | accaacttca | agcacgtggc | ggggcgagca | 540  |
| gcggcggtg  | ctgtgtggg  | gggcttggg  | ggctacgcca | tggggcgct  | tatgtcagg  | 600  |
| atgaactacc | acttcgatag | acccgatgag | taccgatggt | ggagtgaaga | ctcggcgct  | 660  |
| tatccaacc  | gggtttacta | ccgggattac | agcagccccg | tgccacagga | cgctctctg  | 720  |
| gccgattgct | ttaacatcac | agtgactgag | tacagcattg | gccctgctgc | caagaagaac | 780  |
| acctccgagg | ctgtggcggc | agcaaaccaa | acggaggtgg | agatggagaa | caaagtggg  | 840  |
| acgaaggta  | tccgcgagat | gtgcgtgcag | cagtaccgcg | agtaccgcct | ggcctcggg  | 900  |
| atccagctgc | accctgctga | caccctggct | gccctcctcc | tctcctcct  | caccacctc  | 960  |
| tttgccatgc | actgatggga | tgcctgccc  | cgccctgtg  | gcagtgaagt | gacatcggt  | 1020 |
| ccccgtgcc  | acccatgggg | tgttccttgt | cctcgctttt | gtccatcttt | ggtgaagat  | 1080 |
| tcccccgct  | gcctccccgc | aggctctgat | ttgggcaaat | gggaggggat | tttgtcctgt | 1140 |
| cctggtcgtg | gcaggacggc | tgtgtgtgt  | ggagtgggat | gccccaaaa  | tggccttcac | 1200 |
| cacttcctcc | tcctctcct  | ttctggggcg | gagatatggg | ctcgccagc  | ccttattgtc | 1260 |
| cctgcaagag | cgtatctgaa | aatcctcttt | gctaacaagc | agggttttac | ctaactctgt | 1320 |
| tagccccagt | gacagcagag | cgcctttccc | cagggcacac | caaccccaag | ctgaggtgct | 1380 |
| tggcagccac | acgtcccatg | gaggctgatg | gggtttgggg | cgccccaaag | aacaccttg  | 1440 |
| gctactgagg | tgcaattgta | gctctttaat | ctgccaatcc | caacctacc  | gtgtagatag | 1500 |
| gaactgcctg | ctctgcattt | tgcattgtgc | aaacacctcc | tgccgcagcg | cccccaaat  | 1560 |
| agagtgattt | gggaatagtg | aggctgaagc | cacagcagct | tgggattggg | ctcatcatat | 1620 |
| caatccatga | tgtttgctt  | ccagctgagc | ctcactgccc | ttttatagcc | tgccagagg  | 1680 |
| aaggagcgc  | tgctaaatgc | ccaaaaaggt | aacactgagc | aaaagcttat | ttcaatgtat | 1740 |
| gatagagaa  | gagtgcattc | cgcacagatc | agccatggga | gcatcgtttg | ccatcagccc | 1800 |
| caaaacccaa | aggatgctaa | aaagcagcca | aaggggaatc | aagcacgcag | ggaagcctt  | 1860 |
| gaatcagctc | aactggattg | aaatggcaaa | aggcatgagt | agaacgaacg | gcaaggggat | 1920 |
| gctggagatc | cacctcctgt | gagcaaattg | ttcgatgcag | ccaatggaac | tattgtctct | 1980 |
| tgtgtctcag | ttgtgtgta  | tgtgtacata | ggctgtagca | tatgtaaagt | tacacgtgtc | 2040 |
| aagctgctcg | caccgcgtag | agctaatatg | tatcatgtat | gtgggcactg | aatgccaccg | 2100 |
| ttggccatac | ccaaccgtcc | taaacgattt | tcacgtcgct | gtaacttaag | tggagatata | 2160 |
| ctttcagtat | attcagcaaa | aggaattc   |            |            |            | 2188 |

## Mouse PrP cDNA (SEQ ID NO:7; GenBank Accession No. M13685)

|            |            |            |             |            |             |     |
|------------|------------|------------|-------------|------------|-------------|-----|
| aattccttca | gaactgaacc | atttcaaccg | agctgaagca  | ttctgccttc | ctagtggtag  | 60  |
| cagccaatt  | taggagagcc | aagcagacta | tcagtcatca  | tggcgaacct | tggctactgg  | 120 |
| ctgctggccc | tctttgtgac | tatgtggact | gatgtcggcc  | tctgcaaaaa | gcggccaaag  | 180 |
| cctggagggt | ggaacaccgg | tggaaagccg | tatcccgggc  | agggaagccc | tggaggcaac  | 240 |
| cgttaccac  | ctcagggtgg | cacctggggg | cagccccacg  | gtggtggctg | gggacaacct  | 300 |
| catgggggca | gctggggaca | acctcatggt | ggtagtggg   | gtcagcccca | tggcggtgga  | 360 |
| tggggccaag | gaggggtac  | ccataatcag | tggaaacaagc | ccagcaaac  | aaaaaccaac  | 420 |
| ctcaagcatg | tggcaggggc | tgcggcagct | ggggcagtag  | tggggggcct | tgggtggctac | 480 |
| atgctgggga | gcgcctgag  | caggcccatg | atccattttg  | gcaacgactg | ggaggaccgc  | 540 |
| tactaccgtg | aaaacatgta | ccgtaccct  | aaccaagtgt  | actacaggcc | agtggatcag  | 600 |
| tacagcaacc | agaacaactt | cgtgcacgac | tgcgtcaata  | tcaccatcaa | gcagcacacg  | 660 |
| gtcaccacca | ccaccaagg  | ggagaacttc | accgagaccg  | atgtgaagat | gatggagcgc  | 720 |
| tgggtggagc | agatgtgcgt | caccagtag  | cagaaggagt  | cccaggccta | ttacgagcgc  | 780 |
| agaagatcca | gcagcaccgt | gcttttctcc | tccccctctg  | tcatectcct | catctccttc  | 840 |
| ctcatcttcc | tgatcgtagg | atgagggagg | ccttctctgt  | tgttccttcg | cattctctgt  | 900 |
| gtctaggctg | ggggaggggt | tatccacctg | tagctcttcc  | aattgaggtg | gttctcattc  | 960 |

- 11 -

```

ttgcttctct gtgtccccc taggctaata cccctggcac tgatgggccc tgggaaatgt 1020
acagtagacc agttgtctct tgcttcaggt ccctttgatg gagtctgtca tcagccagtg 1080
ctaaccacgg gccataaaga atataacacc aaataactgc tggctagttg gggctttgtt 1140
ttggctctagt gaataaatac tgggtgatcc cctgacttgt acccagagta caaggtgaca 1200
gtgacacatg taacttagca taggcaaagg gttctacaac caaagaagcc actgtttggg 1260
gatggcgccc tggaaaacag cctcccacct gggatagcta gagcatccac acgtggaatt 1320
ctttctttac taacaaacga tagctgattg aaggcaacaa aaaaaaaaaa atcaaattgt 1380
cctactgacg ttgaaagcaa acctttgttc attcccaggg cactagaatg atctttagcc 1440
ttgcttggat tgaactagga gatcttgact ctgaggagag ccagccctgt aaaaagcttg 1500
gtcctcctgt gacgggaggg atggttaagg tacaaggct agaaacttga gtttcttcat 1560
ttctgtctca caattatcaa aagctagaat tagcttctgc cctatgtttc tgtacttcta 1620
tttgcaactgg ataacagaga gacaatctaa acattctctt aggctgcaga taagagaaatg 1680
aggctccatt ccaaagtggg aaagaaattc tgctagcatt gtttaaatac ggcaaaattt 1740
gttcctgaag ttgcttttta ccccagcaga cataaactgc gatagcttca gcttgcaactg 1800
tggattttct gtatagaata tataaaacat aacttcaagc ttatgtcttc tttttaaaac 1860
atctgaagta tgggacgccc tggccgttcc atccagtact aaatgcttac cgtgtgaccc 1920
ttgggctttc agcgtgcact cagttccgta ggattccaaa gcagaccctt agctgggtctt 1980
tgaatctgca tgtacttcac gttttctata tttgtaactt tgcatgtatt ttgttttgtc 2040
atataaaaag tttataaatg tttgctatca gactgacatt aaatagaagc tatgatg 2097

```

## Sheep PrP cDNA (SEQ ID NO:8; GenBank Accession No. X79912)

```

gcagagaagt catcatgggtg aaaagccaca taggcagttg gatcctgggt ctctttgtgg 60
ccatgtggag tgacgtgggc ctctgcaaga agcgaccaa acctggcgga ggatggaaca 120
ctggggggag ccgatacccc ggacagggca gtcctggagg caaccgctat ccacctcagg 180
gaggggggtg ctgggggtcag ccccatggag gtggctgggg ccaacctcat ggagggtggc 240
gggggtcagcc ccaggtgggt ggctggggac agccacatgg tggtaggagg tggggtcagg 300
gtggtagcca cagtcagttg aacaagccca gtaagccaaa aaccaacatg aagcatgtgg 360
caggagctgc tgcagctgga gcagtggtag ggggccttgg tggctacatg ctgggaagtg 420
ccatgagcag gcctcttata cattttgcca atgactatga ggaccgttac tatcgtgaaa 480
acatgtaccg ttaccccaac caagtgtact acagaccagt ggatcagtat agtaaccaga 540
acaactttgt gcatgactgt gtcaacatca cagtcaagca acacacagtc accaccacca 600
ccaaggggga gaacttcacc gaaactgaca tcaagataat ggagcgagtg gtggagcaaa 660
tgggtcagcc ccagttaccag agagaatccc aggcatttta ccaaaggggg gcaagtgtga 720
tcctcttttc tccccctcct gtgatcctcc tcactctctt cctcattttt ctcatagtag 780
gataggggca accttctctg ttt 803

```

## Rat PrP cDNA (SEQ ID NO:9; GenBank Accession No. NM\_012631)

```

atggcgaaac ttggctactg gctgctggcc ctctttgtga ctacatgtac tgatgttggc 60
ctctgcaaaa agcggccaaa gcctggaggg tggaaactgc gtggaagccg gtaccctggg 120
caggggaagcc ctggaggcaa ccgttaccca cctcagagtg gtggtacctg ggggcagccc 180
catggtgggtg gctggggaca acctcatggt ggtggctggg gacaacctca tgggtgtggc 240
tgggggtcagc cccatggcgg gggctggagt caaggagggg gtaccataaa tcagtggaaac 300
aagcccagca agccaaaaac caacctcaag catgtggcag gggctgccgc agctggggca 360
gtagtggggg gccttgggtg ctacatgttg gggagtgcga tgagcaggcc catgctccat 420
tttggcaacg actgggagga ccgctactac cgagaaaaca tgtaccgtta ccctaaccaa 480
gtgtactaca ggccggtgga tcagtacagc aaccagaaca acttcgtgca cgactgtgtc 540
aatatcacca tcaagcagca tacagtcacc accaccacca agggggagaa cttcacggag 600
accgacgtga agatgatgga gcgtgtgggt gagcagatgt gcgtcaccca gtatcagaag 660
gagtcacagg cctattacga cgggagaaga tctagcgccg tgcttttctc cccccctcct 720
gtgatcctcc tcactctctt cctcatcttc ctgatcgtgg gatga 765

```

As used herein, "prion disease" refers to any disease or condition in a subject, the pathogenesis of which involves a prion protein other than PrP<sup>C</sup> of the species of the subject.

- 12 -

A prion disease will typically but not necessarily be a transmissible spongiform encephalopathy.

As used herein, “transmissible spongiform encephalopathy” and, equivalently, “(TSE)” shall mean any prion disease that is associated with spongiform encephalopathy and is communicable from one individual to another. As prion diseases can include entities other than TSE, this term refers to at least a subset of all prion diseases. At present TSE includes Creutzfeldt-Jakob disease, kuru, Gerstmann-Sträussler-Scheinker syndrome, fatal familial insomnia, bovine spongiform encephalopathy, and scrapie.

As used herein, “Creutzfeldt-Jakob disease” and, equivalently, “(CJD)” refers to the TSE that naturally occurs in humans. CJD includes sporadic, genetic (familial), and infectious (i.e., variant and iatrogenic) forms. CJD is a well described entity in the medical literature, and until now has been widely believed to be a uniformly fatal neurodegenerative disease for which there is no effective form of treatment.

As used herein, “variant Creutzfeldt-Jakob disease” and, equivalently, “(vCJD)”, also referred to in the literature as new variant Creutzfeldt-Jakob disease (nvCJD), refers to CJD attributable to the BSE prion. It can be distinguished from sporadic CJD not only by the prion involved but also by certain clinical and preclinical features. See Aguzzi A (2000) *Haematologica* 85:3-10; Hill AF et al. (1997) *Nature* 389:448-50; Bruce ME et al. (1997) *Nature* 389:498-501; Will R et al. (1996) *Lancet* 347:921-5.

As used herein, “iatrogenic Creutzfeldt-Jakob disease” and, equivalently, “(iCJD)” refers to any form of CJD that is attributable to work- or treatment-related exposure to prion protein that is associated with CJD.

As used herein, “bovine spongiform encephalopathy” and, equivalently, “(BSE)” shall refer to the TSE that occurs naturally in cows and cattle.

As used herein, “scrapie” refers to the TSE that occurs naturally in sheep and goats, as well as to experimental models of scrapie. Scrapie in sheep has been recognized and described in the literature for over 300 years. For a review, see O’Rourke KI (2001) *Vet Clin North Am Food Anim Pract* 17:283-300.

As used herein, a “prion protein that is associated with a prion disease” refers to any prion protein involved in the pathogenesis of a prion disease. A prion protein that is associated with a prion disease can be a prion found in nature. Alternatively, prion protein that is associated with a prion disease can be a prion protein made de novo or modified from

- 13 -

its natural form through human activity, e.g., by in vitro synthesis, chemical synthesis, chemical derivativization, or genetic alteration. Chemical alteration includes, without limitation, altered glycosylation. In a preferred embodiment, a prion protein that is associated with a prion disease is a prion protein found in nature, e.g., PrP<sup>Sc</sup>. In one embodiment, a prion protein that is associated with a prion disease is a truncated or genetically modified form of a prion protein found in nature. A genetically modified form of a prion protein found in nature includes a fusion protein involving at least a substantial portion of a prion protein as one component, a prion protein that differs from a prion protein found in nature by one or more conservative amino acid substitutions, and allelic variants of the prion protein found in nature.

Naturally occurring residues can be divided into the following classes based on common side chain properties: (1) hydrophobic: norleucine, Met, Ala, Val, Leu, Ile; (2) neutral hydrophilic: Cys, Ser, Thr; (3) acidic: Asp, Glu; (4) basic: Asn, Gln, His, Lys, Arg; (5) residues that influence chain orientation: Gly, Pro; and (6) aromatic: Trp, Tyr, Phe. Thus, for example, conservative amino acid substitutions can involve the exchange of a member from one of these classes for another member from the same class. Non-conservative amino acid substitutions can involve the exchange of a member of one of these classes for a member from another class.

A conservative amino acid substitution can involve a substitution of a native amino acid residue with another residue such that there is little or no effect on the polarity or charge of the amino acid residue at that position. Conservative amino acid substitutions also encompass non-naturally occurring amino acid residues that are typically incorporated by chemical peptide synthesis rather than by synthesis in biological systems. These include peptidomimetics, and other reversed or inverted forms of amino acid moieties.

As used herein, "prion protein:scrapie form (PrP<sup>Sc</sup>)" refers to any of a number of naturally occurring, species-specific, proteinase K-resistant forms of prion protein associated with prion disease. Specific examples include, but are not limited to, the following: human PrP<sup>Sc</sup> having an amino acid sequence provided by SEQ ID NO:10; bovine PrP<sup>Sc</sup> having an amino acid sequence provided by SEQ ID NO:11; bovine PrP<sup>Sc</sup> having an amino acid sequence provided by SEQ ID NO:12; ovine PrP<sup>Sc</sup> having an amino acid sequence provided by SEQ ID NO:13; ovine PrP<sup>Sc</sup> having an amino acid sequence provided by SEQ ID NO:14; and murine PrP<sup>Sc</sup> having an amino acid sequence provided by SEQ ID NO:15.

- 14 -

Human PrP<sup>Sc</sup> (SEQ ID NO:10; GenBank Accession No. AAE81600)

|   |     |
|---|-----|
| MANLGCWMLV L FVATWSDLG LCKKRPKPGG WNTGGSRYPG QGSPGGNRY P QGGGGWGQP    | 60  |
| HGGGGWGQPHG GGGGGWGQPHG GGGGGWGQPHG GGGGGTHSQWN KPSKPKTNMK HMAGAAAAGA | 120 |
| VVGGLGGGYML GSAMSRPIIH FGSDYEDRY RENMHRYPNQ VYYRPMDEYS NQNNFVHDCV     | 180 |
| NITIKQHTVT TTTKGENFTE TDVKMMERVV EQMCITQYER ESQAYYQRGS SMVLFSSPPV     | 240 |
| ILLISFLIFL IVG  | 253 |

Bovine PrP<sup>Sc</sup> (SEQ ID NO:11; GenBank Accession No. AAE81601)

|  |     |
|--|-----|
| MVKSHIGSWI LVL FVAMWSD VGLCKKRPKP GGWNTGGSRY PGQGSPPGGR YPPQGGGGWG | 60  |
| QPHGGGGWGQ HGGGGWGQPHG GGGGGWGQPHG GGGGGTHGQWN KPSKPKTNMK          | 120 |
| HVAGAAAAGA VVGGLGGGYML GSAMSRPLIH FGSDYEDRY RENMHRYPNQ VYYRVPDQYS  | 180 |
| NQNNFVHDCV NITVKEHTVT TTTKGENFTE TDIKMMERVV EQMCVTQYQK ESQAYYDQGA  | 240 |
| SVILFSSPPV ILLISFLIFL IVG  | 263 |

Bovine PrP<sup>Sc</sup> (SEQ ID NO:12; GenBank Accession No. CAA39368)

|  |     |
|--|-----|
| MVKSHIGSWI LVL FVAMWSD VGLCKKRPKP GGGWNTGGSRY YPGQGSPPGGR YPPQGGGGWG | 60  |
| GQPHGGGGWGQ PHGGGGWGQPH GGGGGWGQPHG GGGGGTHGQWN NKPSKPKTNM           | 120 |
| KHVAGAAAAG AVVGGLGGGYM LGSAMSRPLI HFGSDYEDRY YRENMHRYPN QVYYRVPDQY   | 180 |
| SNQNNFVHDC VNITVKEHTV TTTTKGENFT ETDIKMMERV VEQMCITQYQ RESQAYYQRG    | 240 |
| ASVILFSSPP VILLISFLIF LIVG   | 264 |

Ovine PrP<sup>Sc</sup> (SEQ ID NO:13; GenBank Accession No. AAE81602)

|  |     |
|--|-----|
| MVKSHIGSWI LVL FVAMWSD VGLCKKRPKP GGWNTGGSRY PGQGSPPGGR YPPQGGGGWG | 60  |
| QPHGGGGWGQ HGGGGWGQPHG GGGGGWGQPHG GGGGGTHGQWN NKPSKPKTNM          | 120 |
| GAVVGGLGGY MLGSAMSRPL IHFGNDYEDR YYRENMYRYP NQVYYRVPDQ YSNQNNFVHD  | 180 |
| CVNITVKQHT VTTTTKGENF TETDIKIMER VVEQMCITQY QRESQAYYQR GASVILFSSP  | 240 |
| PVILLISFLI FLIVG   | 255 |

Ovine PrP<sup>Sc</sup> (SEQ ID NO:14; GenBank Accession No. CAA56283)

|  |     |
|--|-----|
| MVKSHIGSWI LVL FVAMWSD VGLCKKRPKP GGGWNTGGSRY YPGQGSPPGGR YPPQGGGGWG | 60  |
| GQPHGGGGWGQ PHGGGGWGQPH GGGGGWGQPHG GGGGGTHGQWN NKPSKPKTNM           | 120 |
| AGAVVGGLGG YMLGSAMSRP LIHFGNDYED RYYRENMYRY PNQVYYRVPD QYSNQN NFVH   | 180 |
| DCVNITVKQH TVTTTTKGEN FTETDIKIME RVVEQMCITQ YQRESQAYYQ RGASVILFSS    | 240 |
| PPVILLISFL IFLIVG  | 256 |

Murine PrP<sup>Sc</sup> (SEQ ID NO:15; GenBank Accession No. AAE81599)

- 15 -

|            |            |            |            |            |            |     |
|------------|------------|------------|------------|------------|------------|-----|
| MANLGYWLLA | LFVTMTDVG  | LCKKRPKPGG | WNTGGSRYPG | QGSPGGNRYP | PQGGTWGQPH | 60  |
| GGGWGQPHGG | SWGQPHGGSW | GQPHGGGWGQ | GGGTHNQWNK | PSKPKTNLKH | VAGAAAAGAV | 120 |
| VGGLGGYMLG | SAMSRPMIHF | GNDWEDRYR  | ENMYRYPNQV | YYRPVDQYSN | QNNFVHDCVN | 180 |
| ITIKQHTVTT | TTKGENFTET | DVKMMERVVE | QMCVTQYQKE | SQAYYDGRRS | SSTVLFSSPP | 240 |
| VILLISFLIF | LIVG       |            |            |            |            | 254 |

As used herein, "full-length PrP<sup>Sc</sup>" refers to a form of PrP<sup>Sc</sup> that includes all its amino acids as it occurs in nature. It is to be distinguished, for example, from a truncated form of PrP<sup>Sc</sup>, described elsewhere herein. A full-length PrP<sup>Sc</sup> can, however, be incorporated into a PrP<sup>Sc</sup> conjugate or PrP<sup>Sc</sup> fusion protein.

As used herein, a "derivative of PrP<sup>Sc</sup>" refers to a chemical or genetic derivative of a naturally occurring form of PrP<sup>Sc</sup>, including PrP<sup>Sc</sup> with non-native glycosylation, covalent or non-covalent conjugates formed between PrP<sup>Sc</sup> and another compound, PrP<sup>Sc</sup> fusion proteins, and any combination thereof. A "derivative of a fragment of PrP<sup>Sc</sup> lacking at least the amino terminus of full-length PrP<sup>Sc</sup>" refers to a chemical or genetic derivative of an N-terminally truncated form of PrP<sup>Sc</sup>, including such truncated forms of PrP<sup>Sc</sup>: (i) with non-native glycosylation, (ii) as part of a covalent or non-covalent conjugate formed with another compound, (iii) as part of a fusion protein, or (iv) any combination thereof. In preferred embodiments the fragment of PrP<sup>Sc</sup> lacking at least the amino terminus of full-length PrP<sup>Sc</sup> refers to a fragment lacking one or more, up to and including all, of the octarepeats (e.g., GGGWGQPH (SEQ ID NO:16) and GGSWGQPH (SEQ ID NO:17)). See Flechsig E et al. (2000) *Neuron* 27:399-408.

A truncated form of a prion protein found in nature is identical in primary sequence to the prion protein found in nature except for the absence of one or more amino acid residues from the N-terminal end, the C-terminal end, or both the N-terminal and the C-terminal ends. Truncated forms can also include deletion mutants, in which an internal sequence is omitted without changing either the N-terminal end or the C-terminal end. In a preferred embodiment, a truncated prion protein lacks one or more, up to and including all, N-terminal octarepeats (e.g., GGGWGQPH and GGSWGQPH). See Flechsig E et al. (2000) *Neuron* 27:399-408.

As used herein, a "fragment of PrP<sup>Sc</sup> lacking at least the amino terminus of full-length PrP<sup>Sc</sup>" shall refer to a truncated form of full-length PrP<sup>Sc</sup> lacking one or more N-terminal amino acids normally present in full-length PrP<sup>Sc</sup>. In a preferred embodiment, the fragment



- 16 -

lacks one or more, up to and including all, of the octarepeats (e.g., GGGWGQPH and GGSWGQPH). See Flechsig E et al. (2000) *Neuron* 27:399-408.

The methods of the instant invention employ immunostimulatory nucleic acids. In the preferred embodiment, the immunostimulatory nucleic acid is a CpG nucleic acid. The terms "nucleic acid" and "oligonucleotide" are used interchangeably to mean multiple nucleotides (i.e., molecules comprising a sugar (e.g., ribose or deoxyribose) linked to a phosphate group and to an exchangeable organic base, which is either a substituted pyrimidine (e.g., cytosine (C), thymidine (T) or uracil (U)) or a substituted purine (e.g., adenine (A) or guanine (G)). As used herein, the terms "nucleic acid" and "oligonucleotide" refer to oligoribonucleotides as well as oligodeoxyribonucleotides. The terms "nucleic acid" and "oligonucleotide" shall also include polynucleosides (i.e., a polynucleotide minus the phosphate) and any other organic base containing polymer. Nucleic acid molecules can be obtained from existing nucleic acid sources (e.g., genomic or cDNA), but are preferably synthetic (e.g., produced by nucleic acid synthesis).

The terms "nucleic acid" and "oligonucleotide" also encompass nucleic acids or oligonucleotides with substitutions or modifications, such as in the bases and/or sugars. For example, they include nucleic acids having backbone sugars that are covalently attached to low molecular weight organic groups other than a hydroxyl group at the 2' position and other than a phosphate group at the 5' position. Thus modified nucleic acids may include a 2'-O-alkylated ribose group. In addition, modified nucleic acids may include sugars such as arabinose instead of ribose. Thus the nucleic acids may be heterogeneous in backbone composition thereby containing any possible combination of polymer units linked together such as peptide-nucleic acids (which have an amino acid backbone with nucleic acid bases).

Nucleic acids also include substituted purines and pyrimidines such as C-5 propyne modified bases. Wagner RW et al. (1996) *Nat Biotechnol* 14:840-4. Purines and pyrimidines include but are not limited to adenine, cytosine, guanine, thymidine, 5-methylcytosine, 2-aminopurine, 2-amino-6-chloropurine, 2,6-diaminopurine, hypoxanthine, and other naturally and non-naturally occurring nucleobases, substituted and unsubstituted aromatic moieties. Other such modifications are well known to those of skill in the art.

The immunostimulatory oligonucleotides of the instant invention can encompass various chemical modifications and substitutions, in comparison to natural RNA and DNA, involving a phosphodiester internucleoside bridge, a  $\beta$ -D-ribose unit and/or a natural

nucleoside base (adenine, guanine, cytosine, thymine, uracil). Examples of chemical modifications are known to the skilled person and are described, for example, in Uhlmann E et al. (1990) *Chem Rev* 90:543; "Protocols for Oligonucleotides and Analogs" Synthesis and Properties & Synthesis and Analytical Techniques, S. Agrawal, Ed, Humana Press, Totowa, USA 1993; Crooke ST et al. (1996) *Annu Rev Pharmacol Toxicol* 36:107-129; and Hunziker J et al. (1995) *Mod Synth Methods* 7:331-417. An oligonucleotide according to the invention can have one or more modifications, wherein each modification is located at a particular phosphodiester internucleoside bridge and/or at a particular  $\beta$ -D-ribose unit and/or at a particular natural nucleoside base position in comparison to an oligonucleotide of the same sequence which is composed of natural DNA or RNA.

For example, the invention relates to an oligonucleotide which comprises one or more modifications and wherein each modification is independently selected from:

- a) the replacement of a phosphodiester internucleoside bridge located at the 3' and/or the 5' end of a nucleoside by a modified internucleoside bridge,
- b) the replacement of phosphodiester bridge located at the 3' and/or the 5' end of a nucleoside by a dephospho bridge,
- c) the replacement of a sugar phosphate unit from the sugar phosphate backbone by another unit,
- d) the replacement of a  $\beta$ -D-ribose unit by a modified sugar unit, and
- e) the replacement of a natural nucleoside base by a modified nucleoside base.

More detailed examples for the chemical modification of an oligonucleotide are as follows.

A phosphodiester internucleoside bridge located at the 3' and/or the 5' end of a nucleoside can be replaced by a modified internucleoside bridge, wherein the modified internucleoside bridge is for example selected from phosphorothioate, phosphorodithioate,  $\text{NR}^1\text{R}^2$ -phosphoramidate, boranophosphate,  $\alpha$ -hydroxybenzyl phosphonate, phosphate-( $\text{C}_1$ - $\text{C}_{21}$ )-O-alkyl ester, phosphate-[( $\text{C}_6$ - $\text{C}_{12}$ )aryl-( $\text{C}_1$ - $\text{C}_{21}$ )-O-alkyl]ester, ( $\text{C}_1$ - $\text{C}_8$ )alkyl-phosphonate and/or ( $\text{C}_6$ - $\text{C}_{12}$ )-arylphosphonate bridges, ( $\text{C}_7$ - $\text{C}_{12}$ )- $\alpha$ -hydroxymethyl-aryl (e.g., disclosed in WO 95/01363), wherein ( $\text{C}_6$ - $\text{C}_{12}$ )aryl, ( $\text{C}_6$ - $\text{C}_{20}$ )aryl and ( $\text{C}_6$ - $\text{C}_{14}$ )aryl are optionally substituted by halogen, alkyl, alkoxy, nitro, cyano, and where  $\text{R}^1$  and  $\text{R}^2$  are, independently of each other, hydrogen, ( $\text{C}_1$ - $\text{C}_{18}$ )-alkyl, ( $\text{C}_6$ - $\text{C}_{20}$ )-aryl, ( $\text{C}_6$ - $\text{C}_{14}$ )-aryl-( $\text{C}_1$ - $\text{C}_8$ )-alkyl, preferably hydrogen, ( $\text{C}_1$ - $\text{C}_8$ )-alkyl, preferably ( $\text{C}_1$ - $\text{C}_4$ )-alkyl and/or methoxyethyl, or  $\text{R}^1$  and  $\text{R}^2$  form, together with

- 18 -

the nitrogen atom carrying them, a 5-6-membered heterocyclic ring which can additionally contain a further heteroatom from the group O, S and N.

The replacement of a phosphodiester bridge located at the 3' and/or the 5' end of a nucleoside by a dephospho bridge (dephospho bridges are described, for example, in Uhlmann E and Peyman A in "Methods in Molecular Biology", Vol. 20, "Protocols for Oligonucleotides and Analogs", S. Agrawal, Ed., Humana Press, Totowa 1993, Chapter 16, pp. 355ff), wherein a dephospho bridge is for example selected from the dephospho bridges formacetal, 3'-thioformacetal, methylhydroxylamine, oxime, methylenedimethyl-hydrazo, dimethylenesulfone and/or silyl groups.

A sugar phosphate unit (i.e., a  $\beta$ -D-ribose and phosphodiester internucleoside bridge together forming a sugar phosphate unit) from the sugar phosphate backbone (i.e., a sugar phosphate backbone is composed of sugar phosphate units) can be replaced by another unit, wherein the other unit is for example suitable to build up a "morpholino-derivative" oligomer (as described, for example, in Stirchak EP et al. (1989) *Nucleic Acids Res* 17:6129-41), that is, e.g., the replacement by a morpholino-derivative unit; or to build up a polyamide nucleic acid ("PNA"; as described for example, in Nielsen PE et al. (1994) *Bioconjug Chem* 5:3-7), that is, e.g., the replacement by a PNA backbone unit, e.g., by 2-aminoethylglycine.

A  $\beta$ -ribose unit or a  $\beta$ -D-2'-deoxyribose unit can be replaced by a modified sugar unit, wherein the modified sugar unit is for example selected from  $\beta$ -D-ribose,  $\alpha$ -D-2'-deoxyribose, L-2'-deoxyribose, 2'-F-2'-deoxyribose, 2'-O-(C<sub>1</sub>-C<sub>6</sub>)alkyl-ribose, preferably 2'-O-(C<sub>1</sub>-C<sub>6</sub>)alkyl-ribose is 2'-O-methylribose, 2'-O-(C<sub>2</sub>-C<sub>6</sub>)alkenyl-ribose, 2'-[O-(C<sub>1</sub>-C<sub>6</sub>)alkyl-O-(C<sub>1</sub>-C<sub>6</sub>)alkyl]-ribose, 2'-NH<sub>2</sub>-2'-deoxyribose,  $\beta$ -D-xylo-furanose,  $\alpha$ -arabinofuranose, 2,4-dideoxy- $\beta$ -D-erythro-hexo-pyranose, and carbocyclic (described, for example, in Froehler J (1992) *Am Chem Soc* 114:8320) and/or open-chain sugar analogs (described, for example, in Vandendriessche et al. (1993) *Tetrahedron* 49:7223) and/or bicyclosugar analogs (described, for example, in Tarkov M et al. (1993) *Helv Chim Acta* 76:481).

A natural nucleoside base can be replaced by a modified nucleoside base, wherein the modified nucleoside base is for example selected from hypoxanthine, uracil, dihydrouracil, pseudouracil, 2-thiouracil, 4-thiouracil, 5-aminouracil, 5-(C<sub>1</sub>-C<sub>6</sub>)-alkyluracil, 5-(C<sub>2</sub>-C<sub>6</sub>)-alkenyluracil, 5-(C<sub>2</sub>-C<sub>6</sub>)-alkynyluracil, 5-(hydroxymethyl)uracil, 5-chlorouracil, 5-fluorouracil, 5-bromouracil, 5-hydroxycytosine, 5-(C<sub>1</sub>-C<sub>6</sub>)-alkylcytosine, 5-(C<sub>2</sub>-C<sub>6</sub>)-alkenylcytosine, 5-(C<sub>2</sub>-C<sub>6</sub>)-alkynylcytosine, 5-chlorocytosine, 5-fluorocytosine,

- 19 -

5-bromocytosine, N<sup>2</sup>-dimethylguanosine, 2,4-diamino-purine, 8-azapurine, a substituted 7-deazapurine, preferably 7-deaza-7-substituted and/or 7-deaza-8-substituted purine or other modifications of a natural nucleoside bases. This list is meant to be exemplary and is not to be interpreted to be limiting.

The oligonucleotides of the present invention are nucleic acids that contain specific sequences found to elicit an immune response. These specific sequences that elicit an immune response are referred to as "immunostimulatory motifs", and the oligonucleotides that contain immunostimulatory motifs are referred to as "immunostimulatory nucleic acid molecules" and, equivalently, "immunostimulatory nucleic acids" or "immunostimulatory oligonucleotides". The immunostimulatory oligonucleotides of the invention thus include at least one immunostimulatory motif.

In one embodiment of the invention the immunostimulatory oligonucleotides include immunostimulatory motifs which are "CpG dinucleotides". A CpG dinucleotide can be methylated or unmethylated. An immunostimulatory nucleic acid containing at least one unmethylated CpG dinucleotide is a nucleic acid molecule which contains an unmethylated cytosine-guanine dinucleotide sequence (i.e., an unmethylated 5' cytosine followed by 3' guanosine and linked by a phosphate bond) and which activates the immune system; such an immunostimulatory nucleic acid is a CpG nucleic acid. CpG nucleic acids have been described in a number of issued patents, published patent applications, and other publications, including U.S. Patent Nos. 6,194,388; 6,207,646; 6,214,806; 6,218,371; 6,239,116; and 6,339,068.

An immunostimulatory nucleic acid containing at least one methylated CpG dinucleotide is a nucleic acid which contains a methylated cytosine-guanine dinucleotide sequence (i.e., a methylated 5' cytosine followed by a 3' guanosine and linked by a phosphate bond) and which activates the immune system. In other embodiments the immunostimulatory oligonucleotides are free of CpG dinucleotides. These oligonucleotides which are free of CpG dinucleotides are referred to as non-CpG oligonucleotides, and they have non-CpG immunostimulatory motifs. The invention, therefore, also encompasses nucleic acids with other types of immunostimulatory motifs, which can be methylated or unmethylated. The immunostimulatory oligonucleotides of the invention, further, can include any combination of methylated and unmethylated CpG and non-CpG

immunostimulatory motifs. In some embodiments the immunostimulatory oligonucleotide is not an antisense oligonucleotide.

As used herein, a "Toll-like receptor (TLR) that signals in response to the CpG nucleic acid" refers to any TLR that engages or initiates an intracellular signaling pathway associated with the development of an immune response, as a result of contacting the TLR with CpG nucleic acid. The pathway typically involves the adaptor protein MyD88 and subsequent downstream molecules including TRAF, IRAK, Jun, Erk, p38 MAPK, and NF- $\kappa$ B. The TLRs are a family of at least ten highly conserved receptors that share as a common feature a cytoplasmic Toll homology IL-1 receptor (TIR) domain believed to be involved in such signaling. TLR9 is reported to be the natural receptor for CpG nucleic acid.

As to CpG nucleic acids, it has recently been described that there are different classes of CpG nucleic acids. One class is potent for activating B cells but is relatively weak in inducing IFN- $\alpha$  and NK cell activation; this class has been termed the B class. The B class CpG nucleic acids typically are fully stabilized and include an unmethylated CpG dinucleotide within certain preferred base contexts. See, e.g., U.S. Patent Nos. 6,194,388; 6,207,646; 6,214,806; 6,218,371; 6,239,116; and 6,339,068. Another class is potent for inducing IFN- $\alpha$  and NK cell activation but is relatively weak at stimulating B cells; this class has been termed the A class. The A class CpG nucleic acids typically have stabilized poly-G sequences at 5' and 3' ends and a palindromic phosphodiester CpG dinucleotide-containing sequence of at least 6 nucleotides. See, for example, published patent application PCT/US00/26527 (WO 01/22990). Yet another class of CpG nucleic acids activates B cells and NK cells and induces IFN- $\alpha$ ; this class has been termed the C class. The C class CpG nucleic acids typically are fully stabilized, include a B class-type sequence and a GC-rich palindrome or near-palindrome. This class has been described in published patent application PCT/US02/26468 (WO 03/015711), the entire content of which is incorporated herein by reference.

Immunostimulatory oligonucleotides are effective in vertebrates. Different immunostimulatory oligonucleotides can cause optimal immune stimulation depending on the type of subject and the sequence of the immunostimulatory oligonucleotide. Many vertebrates have been found according to the invention to be responsive to the same class of immunostimulatory oligonucleotides, sometimes referred to as human specific immunostimulatory oligonucleotides. Rodents, however, respond to different nucleic acids.

- 21 -

Immunostimulatory oligonucleotides causing optimal stimulation in humans may not generally cause optimal stimulation in a mouse and vice versa. An immunostimulatory oligonucleotide causing optimal stimulation in humans often does, however, cause optimal stimulation in other animals such as cow, horses, sheep, etc. For example, within Class B CpG ODN, preferred immunostimulatory sequences have been identified for use in mice (ODN 1826, 5'- TCCATGACGTTTCCTGACGTT -3', SEQ ID NO:18) and for use in humans (ODN 2006, 5'- TCGTCGTTTTGTCGTTTTGTCGTT -3', SEQ ID NO:19). One of skill in the art can identify the optimal immunostimulatory nucleic acid sequences useful for a particular species of interest using routine assays described herein and/or known in the art, using the guidance supplied herein.

As used herein, the term "treat" as used in reference to a disease or condition shall mean to intervene in such disease or condition so as to prevent or slow the development of, slow the progression of, halt the progression of, or eliminate the disease or condition. Thus the phrase "to treat the prion disease" as used herein means to prevent or slow the development of, slow the progression of, halt the progression of, or eliminate the prion disease.

As used herein, a "subject" refers to a human or non-human vertebrate. Preferred non-human vertebrates include feed livestock susceptible to TSE, including cows and cattle, sheep, goats, and pigs. Non-human subjects also specifically include non-human primates as well as rodents. Non-human subjects also include, without limitation, chickens, horses, dogs, cats, guinea pigs, hamsters, mink, and rabbits.

As used herein, a "subject having a prion disease" is a subject known or diagnosed to have a prion disease as disclosed herein. Generally a subject having a prion disease will have some objective manifestation of the prion disease, such as a sign, symptom, or result of a suitable diagnostic test that indicates the presence of a prion disease. In the transmissible spongiform encephalopathies, such objective manifestations can include dementia, ataxia, myoclonus, tremor, presence of protease-resistant prion protein in brain extract, and typical or characteristic abnormalities on brain CT, brain MRI, and/or EEG. This list is not meant to be limiting in any way, and those of skill in the art will recognize what criteria are suitable for making a diagnosis of prion disease in a given species in question. A subject having a prion disease shall also include any subject having a test result which specifically indicates the presence in that subject of any amount of prion protein that is associated with a prion

- 22 -

disease, since it is believed that prion protein that is associated with a prion disease is not present in a subject without a prion disease.

A subject having a prion disease can but need not necessarily have an identifiable risk factor for having a prion disease. An identifiable risk factor for having a prion disease can include a family history of prion disease, a history of consuming known or suspected prion-diseased tissue, or a history of exposure to a prion protein that is associated with a prion disease or to a product derived from a known or suspected prion-diseased tissue (e.g., through administration of pituitary extract).

As used herein, a "subject at risk of developing a prion disease" is a subject with a known or suspected exposure to prion-diseased tissue, a known or suspected exposure to prion protein that is associated with a prion disease, or a known or suspected predisposition to develop a prion disease (e.g., family history of prion disease). In one embodiment the subject at risk of developing a prion disease is a subject residing in or traveling to an area in which TSE is endemic. In one embodiment the subject at risk of developing a prion disease is a subject residing in or traveling to an area in which food or water contains or is likely to contain prion protein that is associated with prion disease. In one embodiment, a subject at risk of developing a prion disease is a subject with a known or suspected iatrogenic exposure to prion-diseased tissue, e.g., neurosurgeons, neuropathologists, pathologists, nurses, morticians, histology technicians and laboratory workers at special risk of contracting iCJD. In one embodiment, a subject at risk of developing a prion disease is a subject with a known or suspected iatrogenic exposure to prion-diseased tissue through receiving a tissue or organ allograft from a subject having a prion disease. Such tissues can include, without limitation, corneas and dural grafts.

As used herein, an "effective amount" of a substance generally refers to that amount of the substance that is sufficient to bring about a desired effect. With reference to CpG nucleic acid, an "effective amount to induce an immune response to the prion protein" shall refer to that amount of CpG nucleic acid that is sufficient to induce an immune response to a particular prion protein. The immune response can occur in vitro, in vivo, ex vivo, and any combination thereof. An immune response to a prion protein can be measured using any suitable means to determine that an immune response occurs in association with exposure of an immune cell to the prion protein. The immune response can be antigen-specific, including any of the following: production of prion protein-specific antibody, proliferation of prion

- 23 -

protein-specific lymphocytes, and cell-mediated immunity against cells expressing prion protein. The immune response can alternatively or additionally be antigen-nonspecific, including any of the following: induction of a Toll-like receptor signaling pathway, inflammation, and production of a cytokine and/or chemokine. Because prion proteins are generally believed not to evoke an immune response, any prion protein-specific immune response which occurs in association with exposure of an immune cell to a prion protein will generally indicate an immune response to the prion protein.

Also with reference to CpG nucleic acid, an "effective amount to treat the prion disease" shall refer to that amount of CpG nucleic acid that is sufficient to treat a particular prion disease. In one embodiment, an effective amount to treat the prion disease is that amount that is sufficient to slow the development of prion disease, compared to the rate of development of prion disease that would occur without CpG administration according to the instant invention. In one embodiment, an effective amount to treat the prion disease is that amount that is sufficient to prevent the development of prion disease, compared to development of prion disease that would occur without CpG administration according to the instant invention. In one embodiment an effective amount to treat the prion disease is that amount that is sufficient to slow the progression of prion disease, compared to the rate of progression of prion disease that would occur without CpG administration according to the instant invention. In one embodiment an effective amount to treat the prion disease is that amount that is sufficient to stop the progression of prion disease, compared to the progression of prion disease that would occur without CpG administration according to the instant invention. In one embodiment an effective amount to treat the prion disease is that amount that is sufficient to resolve prion disease, compared to the prion disease that would occur without CpG administration according to the instant invention.

Combined with the teachings provided herein, by choosing among the various active compounds and weighing factors such as potency, relative bioavailability, patient body weight, severity of adverse side-effects and preferred mode of administration, an effective prophylactic or therapeutic treatment regimen can be planned which does not cause substantial toxicity and yet is entirely effective to treat the particular subject. The effective amount for any particular application can vary depending on such factors as the disease or condition being treated, the particular immunostimulatory oligonucleotide being administered, the antigen, the size of the subject, or the severity of the disease or condition.



- 24 -

One of ordinary skill in the art can empirically determine the effective amount of a particular immunostimulatory oligonucleotide and/or other therapeutic agent without necessitating undue experimentation.

Subject doses of the immunostimulatory oligonucleotides for mucosal or local delivery typically range from about 0.1  $\mu$ g to 10 mg per administration, which depending on the application could be given daily, weekly, or monthly and any other amount of time therebetween. More typically doses range from about 10  $\mu$ g to 5 mg per administration, and most typically from about 100  $\mu$ g to 1 mg, with repeated administrations being spaced days or weeks apart. Subject doses of immunostimulatory oligonucleotides for parenteral delivery for the purpose of inducing an antigen-specific immune response, wherein the compounds are delivered with an antigen but not another therapeutic agent are typically 5 to 10,000 times higher than the effective mucosal dose for vaccine adjuvant or immune stimulant applications, and more typically 10 to 1,000 times higher, and most typically 20 to 100 times higher. Doses of the immunostimulatory oligonucleotides for parenteral delivery for the purpose of inducing an innate immune response or for inducing an antigen-specific immune response when the immunostimulatory nucleic acids are administered in combination with other therapeutic agents or in specialized delivery vehicles typically range from about 0.1  $\mu$ g to 10 mg per administration, which depending on the application could be given daily, weekly, or monthly and any other amount of time therebetween. More typically parenteral doses for these purposes range from about 10  $\mu$ g to 5 mg per administration, and most typically from about 100  $\mu$ g to 1 mg, with repeated administrations being spaced days or weeks apart. In some embodiments, however, parenteral doses for these purposes may be used in a range of 5 to 10,000 times higher than the typical doses described above.

For any immunostimulatory oligonucleotide the therapeutically effective amount can be initially determined from animal models. A therapeutically effective dose can also be determined from human data for CpG oligonucleotides which have been tested in humans (human clinical trials have been initiated) and for compounds which are known to exhibit similar pharmacological activities, such as other mucosal adjuvants, e.g., LT and other antigens for vaccination purposes, for mucosal or local administration. Higher doses are required for parenteral administration. The applied dose can be adjusted based on the relative bioavailability and potency of the administered immunostimulatory oligonucleotide. Adjusting the dose to achieve maximal efficacy based on the methods described herein and

- 25 -

other methods as are well-known in the art is well within the capabilities of the ordinarily skilled artisan.

The immunostimulatory oligonucleotide can be administered alone or with antigen or other therapeutic agent. In this context, "antigen" refers to any biological molecule capable of eliciting specific immunity. Antigens specifically include peptides (oligopeptides, polypeptides, proteins, and glycosylated derivatives thereof), and polysaccharides. Peptide antigen can be administered preformed or as a polynucleotide encoding the peptide. Also in this context, "other therapeutic agent" includes any suitable composition useful in treating prion disease, including an antibody capable of binding a prion protein. When the immunostimulatory oligonucleotide is administered with antigen or other therapeutic agent, the immunostimulatory oligonucleotide can be administered before, concurrently with, or following administration of the antigen or other therapeutic agent. The immunostimulatory oligonucleotide and the antigen or other therapeutic agent can be formulated together or separately when the immunostimulatory oligonucleotide is administered concurrently with the antigen or other therapeutic agent. When the immunostimulatory oligonucleotide is administered before or following administration of antigen or other therapeutic agent, the immunostimulatory oligonucleotide and the antigen or other therapeutic agent can be administered by the same route of administration or by different routes of administration. In addition, when the immunostimulatory oligonucleotide is administered before or following administration of antigen or other therapeutic agent, the immunostimulatory oligonucleotide and the antigen or other therapeutic agent can be administered to the same site or to different sites.

The immunostimulatory oligonucleotides may be directly administered to the subject or may be administered in conjunction with a nucleic acid delivery complex. A nucleic acid delivery complex shall mean a nucleic acid molecule associated with (e.g., ionically or covalently bound to; or encapsulated within) a targeting means (e.g., a molecule that results in higher affinity binding to target cell (e.g., B cell surfaces and/or increased cellular uptake by target cells). Examples of nucleic acid delivery complexes include nucleic acids associated with a sterol (e.g., cholesterol), a lipid (e.g., a cationic lipid, virosome or liposome), or a target cell specific binding agent (e.g., a ligand recognized by target cell specific receptor). Preferred complexes may be sufficiently stable *in vivo* to prevent significant uncoupling prior to internalization by the target cell. However, the complex can

- 26 -

be cleavable under appropriate conditions within the cell so that the nucleic acid is released in a functional form.

Delivery vehicles or delivery devices for delivering antigen and nucleic acids to surfaces have been described. The immunostimulatory oligonucleotide and/or the antigen and/or other therapeutics may be administered alone (e.g., in saline or buffer) or using any delivery vehicles known in the art. For instance the following delivery vehicles have been described: cochleates (Gould-Fogerite et al., 1994, 1996); emulsomes (Vancott et al., 1998, Lowell et al., 1997); ISCOMs (Mowat et al., 1993, Carlsson et al., 1991, Hu et al., 1998, Morein et al., 1999); liposomes (Childers et al., 1999, Michalek et al., 1989, 1992, de Haan 1995a, 1995b); live bacterial vectors (e.g., *Salmonella*, *Escherichia coli*, bacillus Calmette-Guérin, *Shigella*, *Lactobacillus*) (Hone et al., 1996, Pouwels et al., 1998, Chatfield et al., 1993, Stover et al., 1991, Nugent et al., 1998); live viral vectors (e.g., Vaccinia, adenovirus, Herpes Simplex) (Gallichan et al., 1993, 1995, Moss et al., 1996, Nugent et al., 1998, Flexner et al., 1988, Morrow et al., 1999); microspheres (Gupta et al., 1998, Jones et al., 1996, Maloy et al., 1994, Moore et al., 1995, O'Hagan et al., 1994, Eldridge et al., 1989); nucleic acid vaccines (Fynan et al., 1993, Kuklin et al., 1997, Sasaki et al., 1998, Okada et al., 1997, Ishii et al., 1997); polymers (e.g., carboxymethylcellulose, chitosan) (Hamajima et al., 1998, Jabbal-Gill et al., 1998); polymer rings (Wyatt et al., 1998); proteosomes (Vancott et al., 1998, Lowell et al., 1988, 1996, 1997); sodium fluoride (Hashi et al., 1998); transgenic plants (Tacket et al., 1998, Mason et al., 1998, Haq et al., 1995); virosomes (Gluck et al., 1992, Mengiardi et al., 1995, Cryz et al., 1998); virus-like particles (Jiang et al., 1999, Leibl et al., 1998).

The immunostimulatory oligonucleotides may be administered by any means known to the skilled artisan. Routes of administration include but are not limited to oral, mucosal, parenteral, intravenous, intramuscular, intraperitoneal, intranasal, intratracheal, sublingual, subcutaneous, intradermal, inhalation, ocular, vaginal, and rectal.

The immunostimulatory oligonucleotides are administered in pharmaceutically acceptable solutions, which may routinely contain pharmaceutically acceptable concentrations of salt, buffering agents, preservatives, compatible carriers, adjuvants, and optionally other therapeutic ingredients.

Suitable liquid or solid pharmaceutical preparation forms are, for example, aqueous or saline solutions for inhalation, microencapsulated, encochleated, coated onto microscopic

- 27 -

gold particles, contained in liposomes, nebulized, aerosols, pellets for implantation into the skin, or dried onto a sharp object to be scratched into the skin. The pharmaceutical compositions also include granules, powders, tablets, coated tablets, (micro)capsules, suppositories, syrups, emulsions, suspensions, creams, drops or preparations with protracted release of active compounds, in whose preparation excipients and additives and/or auxiliaries such as disintegrants, binders, coating agents, swelling agents, lubricants, flavorings, sweeteners or solubilizers are customarily used as described above. The pharmaceutical compositions are suitable for use in a variety of drug delivery systems. For a brief review of methods for drug delivery, see Langer R (1990) *Science* 249:1527-33, which is incorporated herein by reference.

The immunostimulatory oligonucleotides and optionally other therapeutics and/or antigens may be administered *per se* (neat) or in the form of a pharmaceutically acceptable salt. When used in medicine the salts should be pharmaceutically acceptable, but non-pharmaceutically acceptable salts may conveniently be used to prepare pharmaceutically acceptable salts thereof. Such salts include, but are not limited to, those prepared from the following acids: hydrochloric, hydrobromic, sulphuric, nitric, phosphoric, maleic, acetic, salicylic, p-toluene sulphonic, tartaric, citric, methane sulphonic, formic, malonic, succinic, naphthalene-2-sulphonic, and benzene sulphonic. Also, such salts can be prepared as alkaline metal or alkaline earth salts, such as sodium, potassium or calcium salts of the carboxylic acid group.

Suitable buffering agents include: acetic acid and a salt (1-2% w/v); citric acid and a salt (1-3% w/v); boric acid and a salt (0.5-2.5% w/v); and phosphoric acid and a salt (0.8-2% w/v). Suitable preservatives include benzalkonium chloride (0.003-0.03% w/v); chlorobutanol (0.3-0.9% w/v); parabens (0.01-0.25% w/v) and thimerosal (0.004-0.02% w/v).

For oral administration, the immunostimulatory oligonucleotides can be formulated readily by combining the active compound(s) with pharmaceutically acceptable carriers well known in the art. Such carriers enable the compounds of the invention to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions and the like, for oral ingestion by a subject to be treated. Pharmaceutical preparations for oral use can be obtained as solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients are, in particular, fillers such as sugars, including lactose, sucrose,

- 28 -

mannitol, or sorbitol; cellulose preparations such as, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth, methyl cellulose, hydroxypropylmethyl-cellulose, sodium carboxymethylcellulose, and/or polyvinylpyrrolidone (PVP). If desired, disintegrating agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginate. Optionally the oral formulations may also be formulated in saline or buffers for neutralizing internal acid conditions or may be administered without any carriers.

Dragee cores are provided with suitable coatings. For this purpose, concentrated sugar solutions may be used, which may optionally contain gum arabic, talc, polyvinyl pyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for identification or to characterize different combinations of active compound doses.

Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules can contain the active ingredients in admixture with filler such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In addition, stabilizers may be added. Microspheres formulated for oral administration may also be used. Such microspheres have been well defined in the art. All formulations for oral administration should be in dosages suitable for such administration.

For buccal administration, the compositions may take the form of tablets or lozenges formulated in conventional manner.

For administration by inhalation, the compounds for use according to the present invention may be conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebulizer, with the use of a suitable propellant, e.g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of e.g., gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

The immunostimulatory oligonucleotides, when it is desirable to deliver them systemically, may be formulated for parenteral administration by injection, e.g., by bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form, e.g., in ampoules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents.

Pharmaceutical formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

Alternatively, the active compounds may be in powder form for constitution with a suitable vehicle, e.g., sterile pyrogen-free water, before use.

The immunostimulatory oligonucleotides may also be formulated in rectal or vaginal compositions such as suppositories or retention enemas, e.g., containing conventional suppository bases such as cocoa butter or other glycerides.

In addition to the formulations described previously, the immunostimulatory oligonucleotides may also be formulated as a depot preparation. Such long acting formulations may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

The pharmaceutical compositions also may comprise suitable solid or gel phase carriers or excipients. Examples of such carriers or excipients include but are not limited to calcium carbonate, calcium phosphate, various sugars, starches, cellulose derivatives, gelatin, and polymers such as polyethylene glycols.

The term "pharmaceutically-acceptable carrier" means one or more compatible solid or liquid filler, diluents or encapsulating substances which are suitable for administration to a

- 30 -

human or other vertebrate animal. The term "carrier" denotes an organic or inorganic ingredient, natural or synthetic, with which the active ingredient is combined to facilitate the application. The components of the pharmaceutical compositions also are capable of being commingled with the compounds of the present invention, and with each other, in a manner such that there is no interaction which would substantially impair the desired pharmaceutical efficiency.

The immunostimulatory oligonucleotides useful in the invention may be delivered in mixtures with additional adjuvant(s), other therapeutics, or antigen(s). A mixture may consist of several adjuvants in addition to the immunostimulatory oligonucleotide or several antigens or other therapeutics.

The particular mode selected will depend, of course, upon the particular adjuvants or antigen selected, the particular condition being treated and the dosage required for therapeutic efficacy. The methods of this invention, generally speaking, may be practiced using any mode of administration that is medically acceptable, meaning any mode that produces effective levels of an immune response without causing clinically unacceptable adverse effects. Preferred modes of administration are discussed above.

The immunostimulatory oligonucleotides may conveniently be presented in unit dosage form and may be prepared by any of the methods well known in the art of pharmacy. All methods include the step of bringing the immunostimulatory oligonucleotides into association with a carrier which constitutes one or more accessory ingredients. In general, the compositions are prepared by uniformly and intimately bringing the compounds into association with a liquid carrier, a finely divided solid carrier, or both, and then, if necessary, shaping the product. Liquid dose units are vials or ampoules. Solid dose units are tablets, capsules and suppositories. For treatment of a patient, depending on activity of the compound, manner of administration, purpose of the immunization (i.e., prophylactic or therapeutic), nature and severity of the disorder, age and body weight of the patient, different doses may be necessary. The administration of a given dose can be carried out both by single administration in the form of an individual dose unit or else several smaller dose units. Multiple administration of doses at specific intervals of weeks or months apart is usual for boosting the antigen-specific responses.

Other delivery systems can include time-release, delayed release or sustained release delivery systems. Such systems can avoid repeated administrations of the compounds,

- 31 -

increasing convenience to the subject and the physician. Many types of release delivery systems are available and known to those of ordinary skill in the art. They include polymer base systems such as poly(lactide-glycolide), copolyoxalates, polycaprolactones, polyesteramides, polyorthoesters, polyhydroxybutyric acid, and polyanhydrides. Microcapsules of the foregoing polymers containing drugs are described in, for example, U.S. Patent 5,075,109. Delivery systems also include non-polymer systems that are: lipids including sterols such as cholesterol, cholesterol esters and fatty acids or neutral fats such as mono-di-and tri-glycerides; hydrogel release systems; sylastic systems; peptide based systems; wax coatings; compressed tablets using conventional binders and excipients; partially fused implants; and the like. Specific examples include, but are not limited to: (a) erosional systems in which an agent of the invention is contained in a form within a matrix such as those described in U.S. Patent Nos. 4,452,775, 4,675,189, and 5,736,152, and (b) diffusional systems in which an active component permeates at a controlled rate from a polymer such as described in U.S. Patent Nos. 3,854,480, 5,133,974 and 5,407,686. In addition, pump-based hardware delivery systems can be used, some of which are adapted for implantation.

### Examples

#### Example 1. CpG-ODN Protects Mice from Scrapie Prion.

Groups of 8 mice each were inoculated intraperitoneally with 100  $\mu$ l of 10% brain homogenates from mice terminally ill with RML scrapie prion strain corresponding to an infectious challenge of approximately  $10^4$  LD<sub>50</sub>. The mice received 0.15 pmol (30  $\mu$ l of 5 nM solution) CpG-ODN (oligonucleotide 1826, 5'- TCCATGACGTTCCCTGACGTT -3', SEQ ID NO:18), which has been shown to be a strong inducer of innate immunity and to confer sterile immunity against certain infectious diseases. Sparwasser T et al. (2000) *Eur J Immunol* 30:3591-7. CpG-ODN was administered intraperitoneally at the time of inoculation (0h) with scrapie prions as well as 4 times at intervals of 24h (4 x q 24h); a second group received 0.15 pmol (30  $\mu$ l of 5 nM solution) CpG-ODN 7h after infection (7h) and 4 times at intervals of 24h (4 x q 24h); a third group received CpG-ODN at 7h after infection and subsequently 20 times at intervals of 24h (20 x q 24h). Controls matched for age and sex were given saline instead of CpG-ODN at the identical time intervals. Additional control experiments were also performed in uninfected mice using saline and brain homogenates of



- 32 -

uninfected mice. All animals were observed and scored daily for clinical signs of disease. Scrapie in mice is characterized by ataxia of gait, tremor, difficulty righting from a supine position, and tail rigidity. Occurrence of two of these four symptoms was used as the end point criterion for establishing a clinical diagnosis of scrapie. Western blots of brain homogenates were performed to confirm the diagnosis. All results were analyzed using the Student's t test (Table 1).

Table 1. Mean incubation time in C57BL/6 mice after inoculation with scrapie strain RML and treatment with CpG-ODN.

| Group | Inoculation                          | Treatment | Regimen           | Time to Terminal Disease (d) $\pm$ SD | Attack Rate |
|-------|--------------------------------------|-----------|-------------------|---------------------------------------|-------------|
| 1     | RML                                  | CpG 1826  | 0h and 4 x q 24h  | 253 $\pm$ 4                           | 8/8         |
| 2     | RML                                  | CpG 1826  | 7h and 4 x q 24h  | 250 $\pm$ 6                           | 8/8         |
| 3     | RML                                  | CpG 1826  | 7h and 20 x q 24h | > 330 no disease                      | 0/8         |
| 4     | RML                                  | Saline    | 0h and 4 x q 24h  | 183 $\pm$ 7                           | 8/8         |
| 5     | RML                                  | Saline    | 7h and 4 x q 24h  | 181 $\pm$ 3                           | 8/8         |
| 6     | RML                                  | Saline    | 7h and 20 x q 24h | 181 $\pm$ 3                           | 8/8         |
| 7     | Saline                               | CpG 1826  | 0h and 4 x q 24h  | No disease                            | 0/8         |
| 8     | Saline                               | CpG 1826  | 7h and 4 x q 24h  | No disease                            | 0/8         |
| 9     | Saline                               | CpG 1826  | 7h and 20 x q 24h | No disease                            | 0/8         |
| 10    | Brain homogenate of uninfected mouse | CpG 1826  | 0h and 4 x q 24h  | No disease                            | 0/8         |
| 11    | Brain homogenate of uninfected mouse | CpG 1826  | 7h and 4 x q 24h  | No disease                            | 0/8         |
| 12    | Brain homogenate of uninfected mouse | CpG 1826  | 7h and 20 x q 24h | No disease                            | 0/8         |

Mice infected with the RML strain which received CpG-ODN at the time of inoculation and 7h post-infection as well as 4 times at intervals of 24h showed a dramatic prolongation of survival time compared to control mice with an increase in survival time of 38% in both cases. These differences were highly significant ( $p < 0.0001$ ). The application of CpG-ODN 7h post-inoculation and 20 times at 24h intervals led to disease-free intervals of more than 330 days. All control groups which were not inoculated with the RML strain remained disease-free, and no harmful effect of CpG-ODN application was observed.

- 33 -

These results showed that the application of CpG-ODN at the time of infection and 7h after infection led to a dramatic prolongation of survival time. This effect can be amplified when CpG-ODN are given for a longer period of time. The application of CpG-ODN for 20 times at 24h intervals results in a disease-free interval of >330 days, which indicates the great potential of CpG-ODN for post-exposure prophylaxis of people with exposure to infection. The mechanism of disease prevention remains to be determined, but it seems that the most likely explanation for this effect is a stimulation of TLR-expressing cells of the innate immune system, e.g., macrophages, monocytes, and especially dendritic cells. Sparwasser T et al. (2000) *Eur J Immunol* 30:3591-7. CpG-ODN has been known to induce resistance against other infectious diseases. Zimmermann S et al. (1998) *J Immunol* 160:3627-30. The induction of extreme prolongation of the incubation time or even resistance to prion disease was a surprising finding in the context of a completely different infectious agent.

The findings presented here show that administration of CpG-ODN prolongs the incubation time by 38% and may have the potential to prevent infection after repeated administration, even when high doses of infectivity are administered intraperitoneally. It may therefore be possible to prevent disease after inadvertent iatrogenic exposure with much lower infectious doses administered peripherally.

#### Example 2. Effect of Timing Between Exposure to Prion and Administration of CpG-ODN.

Mice are injected with RML scrapie prion or control and treated with CpG-ODN or control essentially as in Example 1, except that the interval between injection with RML scrapie prion or control and administration of CpG-ODN or control is varied. In some groups administration of CpG-ODN is delayed as much as a month following injection with RML scrapie prion or control. In some groups administration of CpG-ODN precedes injection with RML scrapie prion or control. In some groups the number and schedule of repeated administrations of CpG-ODN is varied from Example 1. Results, measured as in Example 1, show that CpG-ODN is effective even when administered more than 7h after injection with RML scrapie prion.

#### Example 3. Mice Protected from Scrapie Prion by CpG-ODN Develop an Immune Response to Prion.

- 34 -

Mice are injected with RML scrapie prion or control and treated with CpG-ODN or control essentially as in Example 1 or Example 2. At various time points following injection with RML scrapie prion or control and treatment with CpG-ODN or control, tissue or blood samples are obtained and analyzed for prion-specific and prion-nonspecific immune response. Presence of an immune response is determined by suitable method or measurement including, without limitation, antibody titer, enzyme-linked immunosorbent assay (ELISA), flow cytometry, cell proliferation assay, cytotoxicity assay, polymerase chain reaction (PCR) and reverse transcriptase-polymerase chain reaction (RT-PCR), Western immunoblot, Northern blot, and Southern blot. General methods for these types of measurements are standard and are suitably adapted to the specific antigen or stimulus being assayed. For example, ELISA is used to measure production of various secreted products, including antibodies, cytokines and chemokines. Cytokines and chemokines in this example include interleukin (IL)-4, IL-10, IL-6, IL-12, IL-18, interferon (IFN)- $\alpha$ , IFN- $\beta$ , IFN- $\gamma$ , tumor necrosis factor (TNF), and IP-10. Flow cytometry is used to measure cell surface and intracellular proteins, including markers associated with immune cell activation. Markers associated with immune cell activation can vary with cell type but include cluster of differentiation (CD) markers such as CD86, major histocompatibility complex (MHC), inducible cytokine receptors, and certain costimulatory molecules. Results show that CpG-ODN induces an immune response to prion protein.

#### Example 4. Selection of CpG-ODN.

Mice are injected with RML scrapie prion or control and treated with CpG-ODN or control essentially as in Example 1 or Example 2. Various CpG-ODN are compared against ODN 1826 for their effectiveness. Results, measured as in Example 1, show that protection is related to the use of species-optimized CpG-ODN.

#### Example 5. Use of CpG-ODN in Alzheimer's Disease Model.

Mice genetically susceptible to developing Alzheimer's-like disease are administered CpG nucleic acid alone or CpG nucleic acid plus antigen (e.g., amyloid precursor protein or A $\beta$ ), either prior to or following onset of Alzheimer's-like disease. Similar mice are administered appropriate control treatment. Animals are monitored for behavioral and histologic evidence of Alzheimer's-like disease.

- 35 -

The foregoing written specification is considered to be sufficient to enable one skilled in the art to practice the invention. The present invention is not to be limited in scope by examples provided, since the examples are intended as a single illustration of one aspect of the invention and other functionally equivalent embodiments are within the scope of the invention. Various modifications of the invention in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and fall within the scope of the appended claims. The advantages and objects of the invention are not necessarily encompassed by each embodiment of the invention.

All references, patents and patent publications that are recited in this application are incorporated in their entirety herein by reference.

We claim:

- 36 -

### Claims

1. A method for treating a prion disease in a subject, comprising:  
administering to a subject having or at risk of developing a prion disease a CpG nucleic acid in an effective amount to treat the prion disease.
2. The method according to claim 1, wherein the administering follows exposure of the subject to a prion protein that is associated with a prion disease.
3. The method according to claim 1, wherein the prion disease is a transmissible spongiform encephalopathy (TSE).
4. The method according to claim 1, wherein the prion disease is scrapie.
5. The method according to claim 1, wherein the prion disease is bovine spongiform encephalopathy (BSE).
6. The method according to claim 1, wherein the prion disease is variant Creutzfeldt-Jakob disease (vCJD).
7. The method according to claim 1, wherein the prion disease is iatrogenic Creutzfeldt-Jakob disease (iCJD).
8. The method according to claim 1, wherein the subject is a human.
9. A method for inducing an immune response to a prion protein, comprising:  
contacting an antigen-presenting cell (APC) with a prion protein; and  
contacting the APC with a CpG nucleic acid in an effective amount to induce an immune response to the prion protein.
10. The method according to claim 9, wherein the immune response is in vivo.

- 37 -

11. The method according to claim 9, wherein the APC is selected from the group consisting of: a B cell, a dendritic cell, a macrophage, and a monocyte.
12. The method according to claim 9, wherein the APC is a dendritic cell.
13. The method according to claim 9, wherein the APC expresses a Toll-like receptor (TLR) that signals in response to the CpG nucleic acid.
14. The method according to claim 13, wherein the TLR is TLR9.
15. The method according to claim 9, wherein the prion protein is prion protein:scrapie form (PrP<sup>Sc</sup>).
16. The method according to claim 9, wherein the prion protein is a fragment of PrP<sup>Sc</sup> lacking at least the amino terminus of full-length PrP<sup>Sc</sup>.
17. The method according to claim 9, wherein the prion protein is a derivative of PrP<sup>Sc</sup> or a derivative of a fragment of PrP<sup>Sc</sup> lacking at least the amino terminus of full-length PrP<sup>Sc</sup>.
18. The method according to claim 9, wherein the CpG nucleic acid is a Class B CpG nucleic acid.
19. The method according to claim 9, wherein the CpG nucleic acid is a Class A CpG nucleic acid.
20. The method according to claim 9, wherein the CpG nucleic acid is a Class C CpG nucleic acid.
21. The method according to claim 9, wherein the CpG nucleic acid is optimized for use in a species of the subject.

## SEQUENCE LISTING

&lt;110&gt; Coley Pharmaceutical GmbH

&lt;120&gt; USE OF CpG NUCLEIC ACIDS IN PRION DISEASE

&lt;130&gt; C01041.70038

&lt;150&gt; US 60/396,432

&lt;151&gt; 2002-07-17

&lt;160&gt; 19

&lt;170&gt; PatentIn version 3.1

&lt;210&gt; 1

&lt;211&gt; 2415

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 1

```

cggcgcgcgcg agcttctcct ctctctcacga ccgaggcaga gcagtcatta tggcgaacct      60
tggctgctgg atgtctggtc tctttgtggc cacatggagt gacctgggcc tctgcaagaa      120
gcgcccgaag cctggaggat ggaacactgg gggcagccga taccggggc agggcagccc      180
tggaggcaac cgctaccac ctccaggcgcg tgggtggctgg gggcagcctc atggtgggtg      240
ctgggggcag cctcatggtg gtggctgggg gcagcccat ggtgggtggc ggggacagcc      300
tcatggtggt ggctggggtc aaggaggtgg caccacagt cagtggaaaca agccgagtaa      360
gccaaaaacc aacatgaagc acatggctgg tgctgcagca gctggggcag tgggtgggggg      420
ccttggcggc tacatgctgg gaagtgccat gacgagccc atcatacatt tcggcagtga      480
ctatgaggac cgttactatc gtgaaaacat gcaccgttac cccaaccaag tgtactacag      540
gcccattgat gagtacagca accagaacaa ctttgtgcac gactgcgtca atatcacaat      600
caagcagcac acggtcacca caaccaccaa gggggagaac ttcaccgaga ccgacgttaa      660
gatgatggag cgcgtgggtg agcagatgtg tatcaccag tacgagaggg aatctcaggc      720
ctattaccag agaggatcga gcatggtcct cttctcctct ccacctgtga tcctcctgat      780
ctctttcttc atcttctga tagtgggatg aggaaggtct tctgttttc accatctttc      840
taatcttttt ccagcttgag ggaggcggta tccacctgca gcccttttag tgggtggtgc      900
tcaactcttc ttctctcttt gtcccggata ggctaataa tacccttggc actgatgggc      960
actggaaaac atagagtaga cctgagatgc tggtaagcc ccctttgatt gagttcatca     1020
tgagccgttg ctaatgccag gccagtaaaa gtataacagc aaataaccat tggttaatct     1080
ggacttatct ttggacttag tgcaacaggt tgaggctaaa acaaactctc gaacagtctg     1140

```

aaataccttt gcctggatac ctctggctcc ttcagcagct agagctcagt atactaatgc 1200  
 cctatcttag tagagatttc atagctatct agagatatct tccattttta gaaaacccga 1260  
 caacatttct gccaggtttg ttagggaggcc acatgatact tattcaaaaa aatcctagag 1320  
 attcttagct cttgggatgc aggtcagcc cgctggagca tgagctctgt gtgtaccgag 1380  
 aactggggtg atgttttact tttcacagta tgggctacac agcagctgtt caacaagagt 1440  
 aaatattgtc acaacactga acctctggct agaggacata ttcacagtga acataactgt 1500  
 aacatatatg aaaggcttct gggacttgaa atcaaagtgt tgggaatggt gcccttggag 1560  
 gcaacctccc atttttagatg tttaaaggac cctatatgtg gcattccttt ctttaaacta 1620  
 taggtaatta aggcagctga aaagtaaatt gccttctaga cactgaaggc aaatctcctt 1680  
 tgtccattta cctggaaacc agaattgatt tgacatacag gagagctgca gttgtgaaag 1740  
 caccatcatc atagaggatg atgtaattaa aaaatgggtc gtgtgcaaag aaaagaactg 1800  
 cttgcatttc tttatttctg tctcataatt gtcaaaaacc agaattaggt caagttcata 1860  
 gtttctgtaa ttggcttttg aatcaaagaa tagggagaca atctaaaaa tatcttaggt 1920  
 tggagatgac agaaatatga ttgatttgaa gtggaaaaag aaattctgtt aatgttaatt 1980  
 aaagtaaaat tattccctga attgtttgat attgtcacct agcagatatg tattactttt 2040  
 ctgcaatgtt attattggct tgcactttgt gagtatctat gtaaaaatat atatgtatat 2100  
 aaaatatata ttgcatagga cagacttagg agttttgttt agagcagtta acatctgaag 2160  
 tgtctaatac attaaacttt gtaaggctac gaatacttaa tatgtgggaa acccttttgc 2220  
 gtggtcctta ggcttacaat gtgcactgaa tcgtttcatg taagaatcca aagtggacac 2280  
 cattaacagg tctttgaaat atgcatgtac tttatatctt ctatatctgt aactttgcat 2340  
 gttcttggtt tgttatataa aaaaattgta aatgtttaat atctgactga aattaaacga 2400  
 gcgaagatga gcacc 2415

<210> 2

<211> 2446

<212> DNA

<213> Mustela sp.

<400> 2

tcattttgtt ttgttttgtt ttgtttgcag ataagccatc atggtgaaaa gccacatagg 60  
 cagctggctc ctggttctct ttgtggccac atggagtgac attggcttct gcaagaagcg 120  
 gccaaagcct ggaggaggct ggaacactgg ggggagccga taccagggc agggcagtc 180  
 tggaggcaac cgctaccac cccagggtgg tggcggtg ggccagccc acgggggtg 240



ctggggacag ccccacgggg gtggctgggg tcagccccac gggggtggcb ggggacagcc 300  
 gcatggtggc ggtggctggg gtcaaggtgg tgggagccac ggtcagtggg gcaagcccag 360  
 taagcccaaa accaacaatga agcatgtggc gggagccgca gcagccgggg cggtcgtggg 420  
 gggcctgggc ggctacatgc tggggagcgc catgagcagg cccctcattc attttggcaa 480  
 cgactatgag gaccgtact accgtgagaa catgtaccgc taccccaacc aagtgtacta 540  
 caagccggtg gatcagtaca gcaaccagaa caacttcgtg catgactgcg tcaacatcac 600  
 ggtcaagcag cacacggtga ccaccaccac caagggcgag aacttcacgg agaccgacat 660  
 gaagatcatg gagcgcgtgg tggagcagat gtgtgtcacc cagtaccagc gagagtccga 720  
 ggcttactac cagagggggg cgagcgccat cctcttctcg cccctccccg tgatcctcct 780  
 catctcactg ctctattctc tgatagtggg atgaggatgg ccttcccatt ctctccatcg 840  
 tcttcacctt ttacaggttg ggggaggggg tgtctaccta cagccctgta gtgggtgggtg 900  
 ctctattcctg cttctcttta tcaccatag gctaattccc ttggccctga tggccctggg 960  
 aaatgtagag cagacccagg atgctattta ttcaagcccc catgtgttgg agtccttcag 1020  
 gggccaatgc tagtgcaggg ctgagaataa cagcaaatca tcattggttg acctagggct 1080  
 gcttttttgt tgttgttgc tagtgcagct gaccgaggct aaaacaattc tcaaaacagt 1140  
 tttcaaatac ctttgcttgg aaacctctgg ctctgtctgc agctagagct cagtacatta 1200  
 atgtcccatc ttagccgtgt cttcatagca acttggggaa gtttttctcc ccactctaaa 1260  
 agaacgcgat tgcacttccc tgtgcaaaga acatttctgc caaatttgaa aggaggccac 1320  
 atgatattca ttcaaaaagc aaactagaa accctttgct cttggacgca agcccgccct 1380  
 gctaggagca ccaaactggg gcgatgggtt gcattctgcg gcgtgggcta tgcggcagcc 1440  
 gaggtgtcca gcgtaaatat tgatgcgacg ctagacctag gcagaggatg tttgcacagg 1500  
 gaatgaacat aatcaacagt gcgaaaatgc tacaaaaaat cccacactgg ggagcagtgt 1560  
 ccttgagggc aagttttttt ctttttggga catttaaagc ccctatatgt ggcattcctt 1620  
 tctttcgtaa cctaaactat agataattaa ggcagttaaa aattgaactt ccttccaggc 1680  
 cccaagagca aatctttgtt cacttacctg gaaaccagaa tgattttgac acagaggaag 1740  
 gtgcagctgt taaaataacc ctcatcctag aagattgcat catggagaaa acgatccgta 1800  
 gacaaaaatg atcgatttc ttcattgctg tctcgtaatt gacagaaacc agaattatgt 1860  
 caagtccatg tttctataat cagcttttga atcaagaat ggaagtccat ccaaaaaaaaa 1920  
 aaaagaaata ccttaggtca cccatgacag aaatacccat tcagggttaga aaaaaggaat 1980  
 tctgttaact gttatttaag taaggcaaaa ttattgtccg gattgttcga tatcatcagc 2040

tagcagataa attagcattc tgcaatgttc ccggcttgca ctgtgcgggt atttgatgtt 2100  
 aaaaaaatt attatatata ttgtgtatga caaacttaga agtttttgct agaggagtta 2160  
 acatctgata tatctaatac accaccagtt ttggaaggta ctaaataactt aatatgtaga 2220  
 aatccttttg cgtggctctc aggcttacac gtgcactgaa tagttttgta tgatagagcc 2280  
 catgtgggtct tcgaaatatg catgtacttt atattttcta tttttgtaac tgggcatgta 2340  
 cttgtataaa aaatgtataa acattcgaac tcttgactag aattaaacag gaactgagtg 2400  
 tgtcccatgt gtttgcagtg acattcacca ccgcaccctg tgttgg 2446

<210> 3  
 <211> 810  
 <212> DNA  
 <213> *Mesocricetus auratus*

<400> 3  
 tcgaaaatct ccctctttag caatttcttg ctcttagagt ttcagcaatt gctttctcgc 60  
 tccattaggc aacctttcat tttctcacct tccccattat gtaacgggag caatgggttc 120  
 tggaccagtc ttccattaaa gatgattttt atagtcgggtg agcgccgtca gggagtgatg 180  
 acacctgggg gcggttttaa ccgtacaatc ccttaaacca gtctggagcg gtgactcatt 240  
 tccccagggg gaagtggcgc ggccattggt gagcacgacg caagccccgc cccaccacgc 300  
 ccggccccgc cctgctaccc ctctgactc actgccccgc ccgtccccc gcggcgtccg 360  
 agcagcagac cgagaaggca catcgagtcc actcgtcgcg tcgggtggcag gtaagcggct 420  
 tctgaagcct ggccccggga aggggtgctgg agccaggcct cggtgaagcct tcggcttccc 480  
 agagccaagc ccggttact ccggctctcg gggcgtgag gccgcggggc tgaggttgag 540  
 tctggctggg aggtgaccgc gcaccgcag ccgcgcgtct ccttgagggg ccgaacccca 600  
 ggagaggcca ggagccatcc ctctctcccg agcccggtc accccagag tcgctcgggg 660  
 atgggggatg ggggatgggg tggcatcttt tgactgtcgt ttgctgtttt cttctctctt 720  
 tgtaatagct acagcgaaca taattttacc cagggttcca ccgtggtctc gtccgtctc 780  
 ggcatctctc agtccagtac ataccaagg 810

<210> 4  
 <211> 4020  
 <212> DNA  
 <213> *Ovis aries*

<400> 4  
 atggtgaaaa gccacatagg cagttggatc ctggttctct ttgtggccat gtggagtga 60

|  |      |
|--|------|
| gtgggcctct gcaagaagcg accaaaacct ggcggaggat ggaacactgg ggggagccga  | 120  |
| taccggggac agggcagtc tggaggcaac cgctatccac ctgaggagg gggaggctgg    | 180  |
| ggtcagcccc atggagggtg ctggggccaa cctcatggag gtggctgggg tcagcccat   | 240  |
| ggtgggtggct ggggacagcc acatgggtgg ggaggctgg gtcaagggtg tagccacagt  | 300  |
| cagtggaaaca agcccagtaa gccaaaaacc aacatgaagc atgtggcagg agctgctgca | 360  |
| gctggagcag tggtaggggg ccttgggtgg tacatgctgg gaagtgccat gagcaggcct  | 420  |
| cttatacatt ttggcaatga ctatgaggac cgttactatc gtgaaaacat gtaccgttac  | 480  |
| cccaaccaag tgtactacag accagtggat cagtatagta accagaacaa ctttgtgcat  | 540  |
| gactgtgtca acatcacagt caagcaacac acagtcacca ccaccacaa gggggagaac   | 600  |
| ttcaccgaaa ctgacatcaa gataatggag cgagtgggtg agcaaagtgt catcaccag   | 660  |
| taccagagag aatcccaggc ttattaccaa aggggggcaa gtgtgatcct cttttcttcc  | 720  |
| cctcctgtga tcctcctcat ctcttctctc attttctca tagtaggata ggggcaacct   | 780  |
| tcctgttttc attatcttct taatctttgc cagggtgggg gagggagtgt ctacctgcag  | 840  |
| ccctgtagtg gtgggtgtct atttcttctc tctctcttgt tacctgtata ataataacct  | 900  |
| tggcgcttac agcactggga aatgacaagc agacatgaga tgctgtttat tcaagtccca  | 960  |
| ttagctcagt attctaagt cccatcttag cagtgtttt gtagcaattt tctcatttgt    | 1020 |
| ttcaagaaca cctgactaca tttccctttg ggaatagcat ttctgccaa tctggaagga   | 1080 |
| ggccacataa tattcattca aaaaaacaaa actggaaatc cttagtctcat agaccagggg | 1140 |
| tccaccctgt tgagagcatg tgcctctgt ctgcagagaa ctataaagga tattctgcat   | 1200 |
| tttgcagggt acatttgtag gtaacacagc catctattgc atcaagaatg gatattcatg  | 1260 |
| caacctttga cttatgggca gaggacatct tcacaaggaa tgaacataat acaaaggctt  | 1320 |
| ctgagactaa aaaattccaa catatggaag aggtgccctt ggtggcagcc ttccattttg  | 1380 |
| tatgtttaag caccttcaag tgatattcct ttcttttagta acataaagta tagataatta | 1440 |
| aggtacctta attaaactac cttctagaca ctgagagcaa atctgttgtt tatctggaac  | 1500 |
| ccaggatgat tttgacattg cttagggtg tgagagttgg actgtaaaga aagctgagtg   | 1560 |
| ctgaagagtt catgcttttg aactatagtg ttggagaaaa ctcttgagag tcccttggac  | 1620 |
| tgaaggaga tcagtcctga atattcattg gaaggactga tgctgaagct gaaactccaa   | 1680 |
| tactttggtc acctgatggg aagaactgaa ggcaggaggg atgctaggaa agactgaagg  | 1740 |
| caggaggaga aggggacgac agaggatgag atggctagat ggcacatgg actcaatgga   | 1800 |
| catgagctta agtaaactcc aggagtggc aatggacagg gagacctggc gtctgcagt    | 1860 |

ccatggtgtc gcagagtcgg acacgattga gtgactaaat tgaggtgacc cagattttaac 1920  
 atagagaatg cagatacaaa actcatattc atttgattga atcttttcct gaaccagtgc 1980  
 tagtgttgga ctggtaaggg tataacagca tatataggtt atgtgatgaa gagatagtgt 2040  
 acatgaaata tgtgcatttc tttattgctg tcttataatt gtcaaaaaag aaaattaggt 2100  
 ccttggtttc tgtaaaattg acttgaatca aaagggaggg atttaaagaa ataaattaga 2160  
 gatgatagaa atctgatcca ttcagagtag aaaaagaaat tccattactg ttattaaaga 2220  
 aggtaaaatt attccctgaa ttgttcaata ttgtcaccta gcagatagac actattctgt 2280  
 actgttttta ctagcttgca ccttggtgta tcttatgtaa aaacatattt gcatatgaca 2340  
 aactttttct gttagagcaa ttaacatctg aaccaccta tgcattacct gtttttgtaa 2400  
 ggtacttttt gtaaggtagt aaggagatgt gggtttaatc cctaggtcag gtaaattccc 2460  
 tagaggaaga aatggcaacc cactccagta ttcttgccag gaaaatccag tgggcagagg 2520  
 agcctggcag ggtacagtct aagagcatgg ggttgcaaag agtgagacaa gacttgagct 2580  
 actgaacaat aaggacaata aatgctgggt cggctaaaag gttcattagg tttttttct 2640  
 gtaagatggc tctagtagta cttgtcttta tcttcattcg aaacaatttt gttagattgt 2700  
 atgtgacagc tcttgatca gcatgcattt gaaaaaaca tcacaattgg taaatttttg 2760  
 tatagccatc ttactattga agatggaaga aaagaagcaa aattttcagc atatcatgct 2820  
 gtacttattt caagaaagat aaccaaaatg caaaaatgta tttgtgaagt gtatggagaa 2880  
 ggggctgcaa ctgatcaagc ttgtcaaagt agtttgtaaa gtttcgtgct ggagatttct 2940  
 tattggacga tgctccacag ttggatatac cagttgaagt tgatagtgat caaattgaga 3000  
 tattgagaat aatcgatgtt ataccacgcg ggagatagct gacatactca aaatatccaa 3060  
 atagaacctt gaaaaccatt tgcaccatct cagttatgtt aatcactttg atgtttgagt 3120  
 tccacataag caaaaaaaca acaacaaaaa aaaatacaac cttgaccata tttgcgcatg 3180  
 cagttctcta ctgaaatgat tgaaaacact ttgtttttaa aaacagattt tgattaacag 3240  
 tgggtacgat acaataacgt agatggaaga aattgtaggg tgagcaaat gaaccacacc 3300  
 accaaaggcc agtcttcctc taaagaagat gtgtgtatgg tgggattgga aagtaatcct 3360  
 ctattatgga ttcttctgga aaacactgct cctaattaga ccaactgaaa acagcactca 3420  
 acgaaaagca tccagaatta gtcaatagaa aacataatct tccatcagga taacgcaaga 3480  
 ctacatattt ctttgatgac ccagcatggc tggagtttct gattcatctg ttgtattcag 3540  
 acgttgcatc tttggatttt ttccatttat ttcagtctac aaaattatca taatggaaaa 3600

aatttccatt ccctggaaga tgtaaagtgc atctggaaaa tttctttgct caaaaagata 3660  
 aaaagttttg tgaacacaga attatgacgt tgcctgaaaa atggcagaag gtagtggaac 3720  
 aaaagagtga ctatgttggt tggtaaagtt cttagtgaat atgaaaaatg tgtcttttat 3780  
 ttttatttaa acaccaaagg cacattttag caaccaata ctgaatctaa aggaaactct 3840  
 tctgtgtggt gtccttacag tgtgcactga tagtttgtat aagaatccag agtgatatct 3900  
 gaaatacgca tgtgcttata ttttttatat ttgtaacttt gcatgtactt gttttgtggt 3960  
 aaaagtttat aaatatttaa tatctgacta aaattaaaca ggagctaaaa ggagtatctt 4020

<210> 5  
 <211> 795  
 <212> DNA  
 <213> Bos taurus

<400> 5  
 atggtgaaaa gccacatagg cagttggatc ctggttctct ttgtggccat gtggagtgc 60  
 gtgggcctct gcaagaagcg accaaaacct ggaggaggat ggaacactgg ggggagccga 120  
 taccaggac agggcagtc tggaggcaac cgttatccac ctcaggagg ggggtggctgg 180  
 ggtcagcccc atggagggtg ctggggccag cctcatggag gtggctgggg ccagcctcat 240  
 ggaggtggct ggggtcagcc ccatggtggt ggctggggac agccacatgg tgggtggaggc 300  
 tggggtcaag gtggtaccca cgggtcaatgg aacaaacca gtaagccaaa aaccaacatg 360  
 aagcatgtgg caggagctgc tgcagctgga gcagtggtag ggggccttgg tggctacatg 420  
 ctgggaagtg ccatgagcag gcctcttata cattttggca gtgactatga ggaccgttac 480  
 tatcgtgaaa acatgcaccg ttaccccaac caagtgtact acaggccagt ggatcagtat 540  
 agtaaccaga acaactttgt gcatgactgt gtcaacatca cagtcaagga acacacagtc 600  
 accaccacca ccaaggggga gaacttcacc gaaactgaca tcaagatgat ggagcgagtg 660  
 gtggagcaaa tgtgcattac ccagtaccag agagaatccc aggcttatta ccaacgaggg 720  
 gcaagtgtga tcctcttctc tccccctct gtgacctcc tcactctttt cctcattttt 780  
 ctcatagtag gatag 795

<210> 6  
 <211> 2188  
 <212> DNA  
 <213> Gallus gallus

<400> 6  
 gaattccctc ggcagccagc tcctccctct cgctatttat tcctttctcc cccccctacg 60  
 ctggatctgg atcatctcaa gccgagcggg gacggcttct tggatcgctc atacataaat 120

atctgtgagt cagaggaagc aaccaccgac cccaagacct caccgagc catggctagg 180  
 ctctcacca cctgctgcct gctggccctg ctgctcgccg cctgcaccga cgtcgccctc 240  
 tccaagaagg gcaaaggcaa acccagtggt ggggggtggg gcgccgggag ccatcgccag 300  
 ccagctacc ccgcccagcc gggctaccct cataaccag ggtaccccca taaccaggg 360  
 taccctaca accctggcta tcccataac ccggctacc ccagaacctc tggctacccc 420  
 cataaccag gttaccagg ctgggggtcaa ggctacaacc catccagcgg aggaagttac 480  
 cacaaccaga agccatggaa acccccaaa accaacttca agcacgtggc gggggcagca 540  
 gcggcgggtg ctgtgggtggg gggcttgggg ggctacgcca tggggcgcg tatgtcaggg 600  
 atgaactacc acttcgatag acccgatgag taccgatggt ggagtggaa ctcggcgcg 660  
 tatcccaacc gggttacta ccgggattac agcagcccg tgccacagga cgtcttcgtg 720  
 gccgattgct ttaacatcac agtgactgag tacagcattg gccctgctgc caagaagaac 780  
 acctccagg ctgtggcggc agcaaacc aaaggaggtg agatggagaa caaagtggtg 840  
 acgaaggtga tccgcgagat gtgcgtgcag cagtaccgcg agtaccgcct ggcctcgggc 900  
 atccagctgc accctgctga cacctggctc gccgtcctcc tcctcctcct caccaccctt 960  
 tttgccatgc actgatggga tgccgtgccc cggccctgtg gcagtggat gacatcgtgt 1020  
 ccccgctgcc acccatgggg tgttcctgt cctcgtttt gtccatctt ggtgaagatg 1080  
 tcccccgct gcctccccgc aggtctctgat ttgggcaaat gggaggggat tttgtcctgt 1140  
 cctggctgtg gcaggacggc tgctggtggt ggagtgggat gcccaaaaaa tggccttcac 1200  
 cacttcctcc tcctcttcct ttctggggcg gagatatggg ctcgtccagc ccttattgtc 1260  
 cctgcaagag cgtatctgaa aatcctcttt gctaacaagc agggttttac ctaatctgct 1320  
 tagccccagt gacagcagag cgcctttccc cagggcacac caaccccaag ctgaggtgct 1380  
 tggcagccac acgtcccatg gaggtgatg ggttttgggg cgtcccaagc aacaccctgg 1440  
 gctactgagg tgcaattgta gctctttaat ctgccaatcc caaccctacc gtgtagatag 1500  
 gaactgcctg ctctgcattt tgcagtctgc aaacacctcc tgccgcagcg ccccaaaat 1560  
 agagtgattt gggaatagtg aggtgaagc cacagcagct tgggattggg ctcacatat 1620  
 caatccatga tgctttgctt ccagctgagc ctcactgcc ttttatagcc tgcccagagg 1680  
 aaggagcgc tgctaaatgc caaaaagggt aacactgagc aaaagcttat ttcaatgtat 1740  
 gatagagaac gagtgcattc cgcacagatc agccatggga gcatcgttt ccatcagccc 1800  
 caaaacccaa aggatgctaa aatgcagcca aagggaatc aagcacgcag ggaaggactt 1860

gaatcagctc aactggattg aaatggcaaa aggcattgagt agaacgaacg gcaaggggat 1920  
 gctggagatc cacctcctgt gagcaaattg ttogatgcag ccaatggaac tattgcttct 1980  
 tgtgcttcag ttgctgctga tgtgtacata ggctgtagca tatgtaaagt tacacgtgtc 2040  
 aagctgctcg caccgcgtag agctaatatg tatcatgtat gtgggactg aatgccaccg 2100  
 ttggccatac ccaaccgtcc taaacgattt tcacgtcgct gtaacttaag tggagatata 2160  
 ctttcagtat attcagcaaa aggaattc 2188

<210> 7

<211> 2097

<212> DNA

<213> Mus musculus

<400> 7

aattccttca gaactgaacc atttcaaccg agctgaagca ttctgccttc ctagtggtac 60  
 cagtccaatt taggagagcc aagcagacta tcagtcattc tggcgaacct tggctactgg 120  
 ctgctggccc tctttgtgac tatgtggact gatgtcgccc tctgcaaaaa gcggccaaag 180  
 cctggagggt ggaacaccgg tggagccgg tatcccgggc agggagccc tggaggcaac 240  
 cgttaccac ctcagggtgg cacctggggg cagccccacg gtggtggctg gggacaacct 300  
 catgggggca gctggggaca acctcatggt ggtagttggg gtcagcccca tggcgggtgga 360  
 tggggccaag gagggggtac ccataatcag tggacaagc ccagcaaac aaaaaccaac 420  
 ctcaagcatg tggcaggggc tgcggcagct ggggcagtag tggggggcct tgggtggctac 480  
 atgctgggga gcgccgtgag caggcccatg atccattttg gcaacgactg ggaggaccgc 540  
 tactaccgtg aaaacatgta ccgtaccct aaccaagtgt actacaggcc agtggatcag 600  
 tacagcaacc agaacaactt cgtgcacgac tgcgtcaata tcaccatcaa gcagcacacg 660  
 gtcaccacca ccaccaaggg ggagaacttc accgagaccg atgtgaagat gatggagcgc 720  
 gtggtggagc agatgtgctt caccagtagc cagaaggagt ccaggccta ttacgacggg 780  
 agaagatcca gcagcaccgt gcttttctcc tccccctctg tcctcctct catctccttc 840  
 ctcatcttcc tgatcgtggg atgaggagg ccttctctgt tgttccttgc cattctcgtg 900  
 gtctaggctg ggggaggggt tatccacctg tagctctttc aattgaggtg gttctcattc 960  
 ttgcttctct gtgtcccca taggctaata cccctggcac tgatggggcc tgggaaatgt 1020  
 acagtagacc agttgctctt tgcttcaggt ccctttgatg gagtctgtca tcagccagtg 1080  
 ctaacaccgg gccaaataaga atataacacc aaataactgc tggctagttg gggctttgtt 1140  
 ttggtctagt gaataaatac tgggtgtatc cctgacttgt acccagagta caaggtgaca 1200

gtgacacatg taacttagca taggcaaagg gttctacaac caaagaagcc actgtttggg 1260  
 gatggcgccc tggaaaacag cctcccacct gggatagcta gagcatccac acgtggaatt 1320  
 ctttctttac taacaaacga tagctgattg aaggcaacaa aaaaaaaaa atcaaattgt 1380  
 cctactgacg ttgaaagcaa acctttgttc attcccaggg cactagaatg atcttttagcc 1440  
 ttgcttggat tgaactagga gatcttgact ctgaggagag ccagccctgt aaaaagcttg 1500  
 gtccctcctgt gacgggaggg atggttaagg taaaaaggct agaaacttga gtttcttcac 1560  
 ttctgtctca caattatcaa aagctagaat tagcttctgc cctatgtttc tgtacttcta 1620  
 tttgaactgg ataacagaga gacaatctaa acattctctt aggctgcaga taagagaagt 1680  
 aggtccatt ccaaagtggg aaagaaattc tgctagcatt gtttaaatca ggcaaaattt 1740  
 gttcctgaag ttgcttttta cccagcaga cataaactgc gatagcttca gcttgactg 1800  
 tggattttct gtatagaata tataaaacat aacttcaagc ttatgtcttc tttttaaac 1860  
 atctgaagta tgggacgccc tggcgttcc atccagtact aaatgcttac cgtgtgaccc 1920  
 ttgggcttcc agcgtgcact cagtccgta ggattccaaa gcagaccct agctggctct 1980  
 tgaatctgca tgtacttcac gtttctata tttgtaactt tgcattgatt ttgtttgtc 2040  
 atataaaaag tttataaatg tttgctatca gactgacatt aaatagaagc tatgatg 2097

<210> 8  
 <211> 803  
 <212> DNA  
 <213> *Ovis aries*

<400> 8  
 gcagagaagt catcatggtg aaaagccaca taggcagttg gatcctgggt ctctttgtgg 60  
 ccatgtggag tgacgtgggc ctctgcaaga agcgaccaa acctggcgga ggatggaaca 120  
 ctggggggag ccgatacccg ggacagggca gtcctggagg caaccgctat ccacctcagg 180  
 gagggggtgg ctggggtcag ccccatggag gtggctgggg ccaacctcat ggaggtggct 240  
 ggggtcagcc ccatggtggt ggctggggac agccacatgg tggaggaggc tggggtcaag 300  
 gtggtagcca cagtcagtgg aacaagccca gtaagccaaa aaccaacatg aagcatgtgg 360  
 caggagctgc tgcagctgga gcagtggtag ggggccttgg tggctacatg ctgggaagtg 420  
 ccatgagcag gcctcttata ctttttgga atgactatga ggaccgttac tatcgtgaaa 480  
 acatgtaccg ttacccaac caagtgtact acagaccagt ggatcagtat agtaaccaga 540  
 acaactttgt gcatgactgt gtcaacatca cagtcaagca acacacagtc accaccacca 600  
 ccaaggggga gaacttcacc gaaactgaca tcaagataat ggagcgagtg gtggagcaaa 660



tgtgcatcac ccagtaccag agagaatccc aggcttatta ccaaaggggg gcaagtgtga 720  
 tcctcttttc tccccctcct gtgatectcc tcattctctt cctcattttt ctcatagtag 780  
 gataggggca accttcctgt ttt 803

<210> 9  
 <211> 765  
 <212> DNA  
 <213> Rattus norvegicus

<400> 9  
 atggcgaacc ttggctactg gctgctggcc ctctttgtga ctacatgtac tgatgttggc 60  
 ctctgcaaaa agcggccaaa gcctggaggg tggaacactg gtggaagccg gtaccctggg 120  
 cagggaagcc ctggaggcaa ccgttaccca cctcagagtgt gtggtacctg ggggcagccc 180  
 catggtggtg gctggggaca acctcatggt ggtggctggg gacaacctca tgggtggtggc 240  
 tggggtcagc cccatggcgg gggctggagt caaggagggg gtaccataa tcagtggaac 300  
 aagcccagca agccaaaaac caacctcaag catgtggcag gggctgccgc agctggggca 360  
 gtagtggggg gccttggtgg ctacatgttg gggagtgcc a tgagcaggcc catgctccat 420  
 tttggcaacg actgggagga ccgtactac cgagaaaaca tgtaccgtta ccctaacc aa 480  
 gtgtactaca ggccggtgga tcagtacagc aaccagaaca acttcgtgca cgactgtgtc 540  
 aatatcacca tcaagcagca tacagtcacc accaccacca agggggagaa cttcacggag 600  
 accgacgtga agatgatgga gcgtgtggtg gagcagatgt gcgtcaccca gtatcagaag 660  
 gagtcccagg cctattacga cgggagaaga tctagcgccg tgcttttctc ctccccctcct 720  
 gtgatectcc tcattctcct cctcattctt ctgacgtgg gatga 765

<210> 10  
 <211> 253  
 <212> PRT  
 <213> Homo sapiens

<400> 10

Met Ala Asn Leu Gly Cys Trp Met Leu Val Leu Phe Val Ala Thr Trp  
 1 5 10 15

Ser Asp Leu Gly Leu Cys Lys Lys Arg Pro Lys Pro Gly Gly Trp Asn  
 20 25 30

Thr Gly Gly Ser Arg Tyr Pro Gly Gln Gly Ser Pro Gly Gly Asn Arg  
 35 40 45

Tyr Pro Pro Gln Gly Gly Gly Gly Trp Gly Gln Pro His Gly Gly Gly  
 50 55 60  
 Trp Gly Gln Pro His Gly Gly Gly Trp Gly Gln Pro His Gly Gly Gly  
 65 70 75 80  
 Trp Gly Gln Pro His Gly Gly Gly Trp Gly Gln Gly Gly Gly Thr His  
 85 90 95  
 Ser Gln Trp Asn Lys Pro Ser Lys Pro Lys Thr Asn Met Lys His Met  
 100 105 110  
 Ala Gly Ala Ala Ala Ala Gly Ala Val Val Gly Gly Leu Gly Gly Tyr  
 115 120 125  
 Met Leu Gly Ser Ala Met Ser Arg Pro Ile Ile His Phe Gly Ser Asp  
 130 135 140  
 Tyr Glu Asp Arg Tyr Tyr Arg Glu Asn Met His Arg Tyr Pro Asn Gln  
 145 150 155 160  
 Val Tyr Tyr Arg Pro Met Asp Glu Tyr Ser Asn Gln Asn Asn Phe Val  
 165 170 175  
 His Asp Cys Val Asn Ile Thr Ile Lys Gln His Thr Val Thr Thr Thr  
 180 185 190  
 Thr Lys Gly Glu Asn Phe Thr Glu Thr Asp Val Lys Met Met Glu Arg  
 195 200 205  
 Val Val Glu Gln Met Cys Ile Thr Gln Tyr Glu Arg Glu Ser Gln Ala  
 210 215 220  
 Tyr Tyr Gln Arg Gly Ser Ser Met Val Leu Phe Ser Ser Pro Pro Val  
 225 230 235 240  
 Ile Leu Leu Ile Ser Phe Leu Ile Phe Leu Ile Val Gly  
 245 250

<210> 11  
 <211> 263  
 <212> PRT  
 <213> Bos taurus  
 <400> 11

Met Val Lys Ser His Ile Gly Ser Trp Ile Leu Val Leu Phe Val Ala  
 1 5 10 15  
 Met Trp Ser Asp Val Gly Leu Cys Lys Lys Arg Pro Lys Pro Gly Gly  
 20 25 30  
 Trp Asn Thr Gly Gly Ser Arg Tyr Pro Gly Gln Gly Ser Pro Gly Gly  
 35 40 45  
 Asn Arg Tyr Pro Pro Gln Gly Gly Gly Gly Trp Gly Gln Pro His Gly  
 50 55 60  
 Gly Gly Trp Gly Gln Pro His Gly Gly Gly Trp Gly Gln Pro His Gly  
 65 70 75 80  
 Gly Gly Trp Gly Gln Pro His Gly Gly Gly Trp Gly Gln Pro His Gly  
 85 90 95  
 Gly Gly Gly Trp Gly Gln Gly Gly Thr His Gly Gln Trp Asn Lys Pro  
 100 105 110  
 Ser Lys Pro Lys Thr Asn Met Lys His Val Ala Gly Ala Ala Ala Ala  
 115 120 125  
 Gly Ala Val Val Gly Gly Leu Gly Gly Tyr Met Leu Gly Ser Ala Met  
 130 135 140  
 Ser Arg Pro Leu Ile His Phe Gly Ser Asp Tyr Glu Asp Arg Tyr Tyr  
 145 150 155 160  
 Arg Glu Asn Met His Arg Tyr Pro Asn Gln Val Tyr Tyr Arg Pro Val  
 165 170 175  
 Asp Gln Tyr Ser Asn Gln Asn Asn Phe Val His Asp Cys Val Asn Ile  
 180 185 190  
 Thr Val Lys Glu His Thr Val Thr Thr Thr Thr Lys Gly Glu Asn Phe  
 195 200 205  
 Thr Glu Thr Asp Ile Lys Met Met Glu Arg Val Val Glu Gln Met Cys  
 210 215 220  
 Val Thr Gln Tyr Gln Lys Glu Ser Gln Ala Tyr Tyr Asp Gln Gly Ala  
 225 230 235 240

Ser Val Ile Leu Phe Ser Ser Pro Pro Val Ile Leu Leu Ile Ser Phe  
 245 250 255

Leu Ile Phe Leu Ile Val Gly  
 260

<210> 12  
 <211> 264  
 <212> PRT  
 <213> Bos taurus

<400> 12

Met Val Lys Ser His Ile Gly Ser Trp Ile Leu Val Leu Phe Val Ala  
 1 5 10 15

Met Trp Ser Asp Val Gly Leu Cys Lys Lys Arg Pro Lys Pro Gly Gly  
 20 25 30

Gly Trp Asn Thr Gly Gly Ser Arg Tyr Pro Gly Gln Gly Ser Pro Gly  
 35 40 45

Gly Asn Arg Tyr Pro Pro Gln Gly Gly Gly Gly Trp Gly Gln Pro His  
 50 55 60

Gly Gly Gly Trp Gly Gln Pro His Gly Gly Gly Trp Gly Gln Pro His  
 65 70 75 80

Gly Gly Gly Trp Gly Gln Pro His Gly Gly Gly Trp Gly Gln Pro His  
 85 90 95

Gly Gly Gly Gly Trp Gly Gln Gly Gly Thr His Gly Gln Trp Asn Lys  
 100 105 110

Pro Ser Lys Pro Lys Thr Asn Met Lys His Val Ala Gly Ala Ala Ala  
 115 120 125

Ala Gly Ala Val Val Gly Gly Leu Gly Gly Tyr Met Leu Gly Ser Ala  
 130 135 140

Met Ser Arg Pro Leu Ile His Phe Gly Ser Asp Tyr Glu Asp Arg Tyr  
 145 150 155 160

Tyr Arg Glu Asn Met His Arg Tyr Pro Asn Gln Val Tyr Tyr Arg Pro  
 165 170 175

Val Asp Gln Tyr Ser Asn Gln Asn Asn Phe Val His Asp Cys Val Asn  
180 185 190

Ile Thr Val Lys Glu His Thr Val Thr Thr Thr Thr Lys Gly Glu Asn  
195 200 205

Phe Thr Glu Thr Asp Ile Lys Met Met Glu Arg Val Val Glu Gln Met  
210 215 220

Cys Ile Thr Gln Tyr Gln Arg Glu Ser Gln Ala Tyr Tyr Gln Arg Gly  
225 230 235 240

Ala Ser Val Ile Leu Phe Ser Ser Pro Pro Val Ile Leu Leu Ile Ser  
245 250 255

Phe Leu Ile Phe Leu Ile Val Gly  
260

<210> 13  
<211> 255  
<212> PRT  
<213> Ovis aries

<400> 13

Met Val Lys Ser His Ile Gly Ser Trp Ile Leu Val Leu Phe Val Ala  
1 5 10 15

Met Trp Ser Asp Val Gly Leu Cys Lys Lys Arg Pro Lys Pro Gly Gly  
20 25 30

Trp Asn Thr Gly Gly Ser Arg Tyr Pro Gly Gln Gly Ser Pro Gly Gly  
35 40 45

Asn Arg Tyr Pro Pro Gln Gly Gly Gly Gly Trp Gly Gln Pro His Gly  
50 55 60

Gly Gly Trp Gly Gln Pro His Gly Gly Gly Trp Gly Gln Pro His Gly  
65 70 75 80

Gly Ser Trp Gly Gln Pro His Gly Gly Gly Gly Trp Gly Gln Gly Gly  
85 90 95

Ser His Ser Gln Trp Asn Lys Pro Ser Lys Pro Lys Thr Asn Met Lys  
100 105 110

His Val Ala Gly Ala Ala Ala Ala Gly Ala Val Val Gly Gly Leu Gly  
 115 120 125

Gly Tyr Met Leu Gly Ser Ala Met Ser Arg Pro Leu Ile His Phe Gly  
 130 135 140

Asn Asp Tyr Glu Asp Arg Tyr Tyr Arg Glu Asn Met Tyr Arg Tyr Pro  
 145 150 155 160

Asn Gln Val Tyr Tyr Arg Pro Val Asp Gln Tyr Ser Asn Gln Asn Asn  
 165 170 175

Phe Val His Asp Cys Val Asn Ile Thr Val Lys Gln His Thr Val Thr  
 180 185 190

Thr Thr Thr Lys Gly Glu Asn Phe Thr Glu Thr Asp Ile Lys Ile Met  
 195 200 205

Glu Arg Val Val Glu Gln Met Cys Ile Thr Gln Tyr Gln Arg Glu Ser  
 210 215 220

Gln Ala Tyr Tyr Gln Arg Gly Ala Ser Val Ile Leu Phe Ser Ser Pro  
 225 230 235 240

Pro Val Ile Leu Leu Ile Ser Phe Leu Ile Phe Leu Ile Val Gly  
 245 250 255

<210> 14  
 <211> 256  
 <212> PRT  
 <213> Ovis aries

<400> 14

Met Val Lys Ser His Ile Gly Ser Trp Ile Leu Val Leu Phe Val Ala  
 1 5 10 15

Met Trp Ser Asp Val Gly Leu Cys Lys Lys Arg Pro Lys Pro Gly Gly  
 20 25 30

Gly Trp Asn Thr Gly Gly Ser Arg Tyr Pro Gly Gln Gly Ser Pro Gly  
 35 40 45

Gly Asn Arg Tyr Pro Pro Gln Gly Gly Gly Gly Trp Gly Gln Pro His  
 50 55 60

Gly Gly Gly Trp Gly Gln Pro His Gly Gly Gly Trp Gly Gln Pro His  
65 70 75 80

Gly Gly Gly Trp Gly Gln Pro His Gly Gly Gly Gly Trp Gly Gln Gly  
85 90 95

Gly Ser His Ser Gln Trp Asn Lys Pro Ser Lys Pro Lys Thr Asn Met  
100 105 110

Lys His Val Ala Gly Ala Ala Ala Gly Ala Val Val Gly Gly Leu  
115 120 125

Gly Gly Tyr Met Leu Gly Ser Ala Met Ser Arg Pro Leu Ile His Phe  
130 135 140

Gly Asn Asp Tyr Glu Asp Arg Tyr Tyr Arg Glu Asn Met Tyr Arg Tyr  
145 150 155 160

Pro Asn Gln Val Tyr Tyr Arg Pro Val Asp Gln Tyr Ser Asn Gln Asn  
165 170 175

Asn Phe Val His Asp Cys Val Asn Ile Thr Val Lys Gln His Thr Val  
180 185 190

Thr Thr Thr Thr Lys Gly Glu Asn Phe Thr Glu Thr Asp Ile Lys Ile  
195 200 205

Met Glu Arg Val Val Glu Gln Met Cys Ile Thr Gln Tyr Gln Arg Glu  
210 215 220

Ser Gln Ala Tyr Tyr Gln Arg Gly Ala Ser Val Ile Leu Phe Ser Ser  
225 230 235 240

Pro Pro Val Ile Leu Leu Ile Ser Phe Leu Ile Phe Leu Ile Val Gly  
245 250 255

<210> 15

<211> 254

<212> PRT

<213> Mus musculus

<400> 15

Met Ala Asn Leu Gly Tyr Trp Leu Leu Ala Leu Phe Val Thr Met Trp  
1 5 10 15

Thr Asp Val Gly Leu Cys Lys Lys Arg Pro Lys Pro Gly Gly Trp Asn  
 20 25 30

Thr Gly Gly Ser Arg Tyr Pro Gly Gln Gly Ser Pro Gly Gly Asn Arg  
 35 40 45

Tyr Pro Pro Gln Gly Gly Thr Trp Gly Gln Pro His Gly Gly Gly Trp  
 50 55 60

Gly Gln Pro His Gly Gly Ser Trp Gly Gln Pro His Gly Gly Ser Trp  
 65 70 75 80

Gly Gln Pro His Gly Gly Gly Trp Gly Gln Gly Gly Gly Thr His Asn  
 85 90 95

Gln Trp Asn Lys Pro Ser Lys Pro Lys Thr Asn Leu Lys His Val Ala  
 100 105 110

Gly Ala Ala Ala Ala Gly Ala Val Val Gly Gly Leu Gly Gly Tyr Met  
 115 120 125

Leu Gly Ser Ala Met Ser Arg Pro Met Ile His Phe Gly Asn Asp Trp  
 130 135 140

Glu Asp Arg Tyr Tyr Arg Glu Asn Met Tyr Arg Tyr Pro Asn Gln Val  
 145 150 155 160

Tyr Tyr Arg Pro Val Asp Gln Tyr Ser Asn Gln Asn Asn Phe Val His  
 165 170 175

Asp Cys Val Asn Ile Thr Ile Lys Gln His Thr Val Thr Thr Thr Thr  
 180 185 190

Lys Gly Glu Asn Phe Thr Glu Thr Asp Val Lys Met Met Glu Arg Val  
 195 200 205

Val Glu Gln Met Cys Val Thr Gln Tyr Gln Lys Glu Ser Gln Ala Tyr  
 210 215 220

Tyr Asp Gly Arg Arg Ser Ser Ser Thr Val Leu Phe Ser Ser Pro Pro  
 225 230 235 240

Val Ile Leu Leu Ile Ser Phe Leu Ile Phe Leu Ile Val Gly  
 245 250



<210> 16  
<211> 8  
<212> PRT  
<213> Mus musculus

<400> 16

Gly Gly Gly Trp Gly Gln Pro His  
1 5

<210> 17  
<211> 8  
<212> PRT  
<213> Mus musculus

<400> 17

Gly Gly Ser Trp Gly Gln Pro His  
1 5

<210> 18  
<211> 20  
<212> DNA  
<213> Artificial sequence

<220>

<223> Synthetic oligonucleotide

<400> 18  
tccatgacgt tcctgacgtt

20

<210> 19  
<211> 24  
<212> DNA  
<213> Artificial sequence

<220>

<223> Synthetic oligonucleotide

<400> 19  
tcgtcgtttt gtcgttttgc cggt

24